arXiv:1610.05145v1 [math.CT] 17 Oct 2016

Globularily generated double categories II: The free globularily generated double category construction

Juan Orendain

Abstract: We introduce the free globularily generated double category construction. We aim to establish an analogy between the way double categories and globularily generated double categories relate to bicategories. We introduce the notion of decorated bicategory and extend the definition of horizontalization functor to the definition of decorated horizontalization functor. We furnish globularily generated double category construction with the structure of a functor and we prove that this functor is left adjoint to decorated horizontalization functor. We regard this relation as a generalization of relations satisfied by horizontalization functor and trivial double category functor in the theory of categorical structures of second order.

Contents

1 Introduction **1**

2 Preliminaries **4**

3 Free globularily generated double category **16**

4 Canonical double projection **34**

5 Free double functor **47**

6 Freeness **60**

7 References **68**

1 Introduction

There exists, in the mathematical literature, a variety of competing ideas of what a higher order categorical structure should be. This set of ideas reduces, in the case of categorical structures of order 2, to the concepts of bicategory and double category, both types of structures introduced by Ehresmann, in [3] and [4]. These two concepts are related in different ways. The most obvious being that every bicategory can be considered as a 'trivial' double category. Less obviously, every double category admits an adjacent trivial

double category, its horizontal bicategory. These relations admit first order categorical extensions. These relations can be summed up by the following statement: Category adjacent to 3-category of bicategories, bifunctors, lax natural transformations, and deformations is a reflective subcategory of category adjacent to 2-category of double categories, double functors, and double natural transformations, with horizontalization functor as reflector. Second order extensions to this statement are studied in [8].

In [7] the concept of globularily generated double category was introduced. The condition of a double category being globularily generated is considered as a generalization of the condition of a double category being trivial. It was proven that 2-category of globularily generated double categories, double functors, and globularily generated double natural transformations is a strictly 2-reflective subcategory of 2-category of general double categories, double functors and globularily generated double natural transformations. A 2-reflector for this relation was explicitly constructed, the globularily generated piece 2-functor. This relation is regarded as an analogy between the way globularily generated double categories and trivial double categories relate to general double categories. In this analogy globularily generated piece 2-functor plays the role of horizontalization functor.

We aim to establish, in this paper, an analogy between the way globularily generated double categories and general double categories relate to bicategories. We introduce the concepts of decorated bicategory and decorated horizontalization. We regard decorated horizontalization functor as a generalized version of horizontalization functor [8]. We introduce the free globularily generated double category construction and we furnish this construction with the structure of a functor. We prove that free globularily generated double category functor is left adjoint to decorated horizontalization functor, thus establishing an analogy with the fact that horizontalization functor is left adjoint to the trivial double category functor. Decorated horizontalization functor plays the role of horizontalization functor and free globularily generated double category functor plays the role of trivial double category functor in this analogy. We prove that free globularily generated double category contains all the information of every globularily generated verticalization of its base decorated bicategory. We equip every free globularily generated double category with a filtration of its morphism category in analogy with the way vertical filtration of a general globularily generated double category is defined [7]. We now sketch the contents of each of the sections that form this paper.

In section 2 we recall basic concepts related to the theory of categorical structures of order 2. We present relevant examples and establish notational

conventions used in the rest of the paper. We review the concept of globularily generated double category already introduced in [7] and we introduce the concepts of decorated bicategory, decorated bifunctor and decorated horizontalization. We prove that decorated horizontalization functor is faithful when restricted to category of globularily generated double categories. In section 3 we introduce the main construction of the paper. Precisely, we associate to each decorated bicategory a globularily generated double category, its free globularily generated double category. We perform the construction of the globularily generated double category of a decorated bicategory in several steps. We equip free globularily generated double category construction with a filtration reminiscent of vertical filtration associated to any globularily generated double category. We present examples illustrating the fact that this filtration not always coincides with vertical filtration and that the free globularily generated double category construction not always provides a verticalization. In section 4 we introduce the canonical projection construction. We associate a double functor from free globularily generated double category associated to a decorated bicategory to any of its globularily generated verticalizations. We prove that this double functor is uniquely defined by a certain set of properties and that its morphism functor is complete. We interpret this by saying that canonical projection canonically provides every globularily generated verticalization of a decorated bicategory with the structure of a quotient of its free globularily generated double category. In section 5 we generalize the canonical projection construction to the free double functor construction. We replicate the ideas behind the construction presented in section 4 to associate to every decorated bifunctor between decorated bicategories a double functor between corresponding free globularily generated double categories. We call this construction the free double functor construction. We prove that free double functor construction is determined uniquely by a certain set of properties and that it restricts in a particular case to the canonical projection functor. In section 6 we prove that free globularily generated double category construction assembles into a functor and that this functor is left adjoint to decorated horizontalization functor. Precisely, we prove that pair formed by function associating free globularily generated double category to every decorated bicategory and free double functor to every decorated bifunctor is a functor. We prove that collection of canonical projections is a natural transformation and that canonical projection transformation forms part of a counit-unit pair making free globularily generated double category construction functor left adjoint to decorated horizontalization functor. We interpret this by saying that free globularily generated double category is a a free construction with respect

to decorated horizontalization functor. We consider this as evidence of an analogy between free globularily generated double category construction and trivial double category functor.

The exposition of the paper will be elementary for the most part. Exposure to the theories of bicategories and of double categories is assumed. We refer the reader to the standard references in the theory of bicategories [5] and in the theory of double categories [8] Our notation may differ from these references. We establish the notational conventions followed in the rest of the paper in section 2.

2 Preliminaries

In this first section we recall basic concepts related to the theory of categorical structures of second order. We review the defininition of globularily generated double category, and introduce the concepts of decorated bicategory, decorated bifunctor, and decorated horizontalization. We prove that restriction of decorated horizontalization functor to category of globularily generated double categories is faithful. We establish notational conventions used throughout the paper and present relevant examples.

Bicategories

We will understand, for a bicategory, a category weakly enriched over 2-category **Cat** of categories, functors, and natural transformations [6]. Given a bicategory B we will write B_0, B_1, and B_2, for collection of 0-,1-, and 2-cells of B respectively, we will write $\bullet$ and $*$ for vertical and horizontal composition of 2-cells in B and we will write λ, ρ, and A for left and right identity transformations and associator of B respectively. In the case in which horizontal identity transformations and associator of bicategory B are identity natural transformations we will say that bicategory B is a 2-category. We will understand, for a bifunctor, a functor strictly enriched over **Cat**. Given a bifunctor F we will write F_0, F_1, and F_2 for 0-,1-, and 2-cell components of F respectively. Composition of bifunctors is performed component-wise. This composition operation is unital and associative, and thus provides pair formed by collection of bicategories and collection of bifunctors between them with the structure of a category. We write **bCat** for this category. Note that category **bCat** is usually studied as the adjacent category of a 3-category whose 2-cells are lax or oplax natural transformations and whose 3-cells are deformations between transformations. We will

not make use of this additional structure in this paper.

Double categories

We understand, for a double category, a category weakly internal to 2-category **Cat** [6]. Given a double category C we will write C_0 for category of objects of C and C_1 for category of morphisms of C. We will denote by s, t, and i source, target, and identity functors of C and by $*$ composition bifunctor of C. We will, in analogy with the theory of bicategories, write λ and ρ for left and right identity transformations of C and A for associator of C. We say that a 2-morphism in double category C is **globular** if its source and target are vertical identity endomorphisms. The definition of double category that we use [8] requires for components of left and right identity transformations and of associator of double categories to be globular. In the case in which horizontal identity transformations and associator of double category C are identity natural transformations we will say that double category C is strict. Given double categories C and C' we understand for a double functor from C to C', a functor from C to C', strictly internal to **Cat**. We will write F_0 and F_1 for object functor and morphism functor of a double functor F. Composition of double functors is performed component-wise. This composition operation is unital and associative, and thus provides pair formed by collection of double categories and collection of double functors between them with the structure of a category. We denote this category by **dCat**. Note that category **dCat** is usually studied as the adjacent category of a 2-category whose 2-cells are double natural transformations. We will not make use of this structure in this paper.

Trivial double categories and horizontalization

Given a bicategory B, we associate to B a double category $\overline{B}$ whose object category is given by discrete category generated by collection of objects B_0 of B, whose collection of horizontal morphisms is given by collection B_1 of 1-cells of B and whose collection of 2-morphisms is given by collection B_2 of globular morphisms of B. The obvious functors serve as source, target, and identity functors of $\overline{B}$. Horizontal composition bifunctor of $\overline{B}$ is bifunctor generated by horizontal composition operation in B. Left and right identity transformations and associator in $\overline{B}$ are given by transformations induced by left and right identity transformations and associator in B. We call $\overline{B}$ the trivial double category associated to bicategory B. Given a bifunctor F between bicategories B and B' we denote by $\overline{F}$ pair formed by functor from

object category of $\overline{B}$ to object category of $\overline{B}'$, generated by 0-cell function F_0 of F, and pair formed by 1-cell function F_1 of F and 2-cell function F_2 of F. Thus defined pair $\overline{F}$ is a double functor from trivial double category $\overline{B}$ associated to B to trivial double category $\overline{B}'$ associated to B'. We call $\overline{F}$ the trivial double functor associated to bifunctor F. Pair formed by functions associating to every bicategory B its trivial double category $\overline{B}$ and function associating to every bifunctor F its trivial double functor $\overline{F}$ is a functor from category **bCat** to category **dCat**. We call this functor the trivial double category functor.

Now, given a double category C, we associate to C a bicategory HC whose collections of 0-,1-, and 2-cells are collection of objects of C, collection of horizontal morphisms of C, and collection of globular morphisms of C respectively. Vertical and horizontal composition of 2-cells in HC is vertical and horizontal composition of 2-cells in C. Left and right identity transformations and associator of HC are induced by left and right identity transformations and associator of C respectively. We call HC the horizontal bicategory, or simply the horizontalization, of C. Given a double functor F between double categories C and C' we denote by HF triple formed by object function of object functor F_0 of F, object function of morphism functor F_1 of F, and morphism function of morphism functor F_1 of F. Thus defined HF is a bifunctor from horizontal bicategory HC associated to C to horizontal bicategory HC' associated to C'. We call HF the horizontal bifunctor, or the horizontalization, of double functor F. Pair H formed by function associating horizontal bicategory HC to every double category C and function associating horizontal bifunctor HF to every double functor F is a functor from category **dCat** to **bCat**. We call this functor the horizontalization functor and we denote it by H.

Given a bicategory B it is easy to see that equation $H\overline{B} = B$ holds. This relations extends to a functorial setting, that is, horizontalization functor is left inverse to trivial double category functor. Horizontalization functor is not right inverse to trivial double category functor. A weaker relation holds. Horizontalization functor is left adjoint to trivial double category functor. Trivial double category functor is injective and faithful, and thus makes category **bCat** a subcategory of **dCat**. Horizontalization functor is thus a reflector for this inclusion.

Globularily generated double categories

We say that a pair D of categories D_0, D_1 is a sub-double category of a double category C if D_0 is a subcategory of object category C_0 of C, D_1 is a sub-

category of morphism category C_1 of C, source and target functors s, t of C restrict to functors from D_1 to D_0, horizontal identity functor i of C restricts to a functor from D_0 to D_1, horizontal composition bifunctor $*$ restricts to a bifunctor from $D_1 \times_{D_0} D_1$ to D_1, components of left and right identity transformations of C, associated to horizontal morphisms in D_1, are morphisms in D_1, and if components of associator of C, associated to composable triples of horizontal morphisms of C in D_1, are morphisms in D_1. Observe that in this case pair D together with restrictions of structure functors s, t, i, and $*$ of C, left and right identity natural isomorphisms, and associator of C, is itself a double category. Given a collection of sub-double categories D_α, $\alpha \in A$, of a double category C, pair formed by intersection $\cap_{\alpha \in A} D_{\alpha_0}$ of object categories of double categories D_α, $\alpha \in A$, and intersection $\cap_{\alpha \in A} D_{\alpha_1}$ of morphism categories of double categories D_α, $\alpha \in A$, is again a sub-double category of C. Given a collection of 2-morphisms X of double category C, we call the intersection of all complete sub-double categories D, of C, such that collection X is contained in collection of 2-morphisms of D, the complete sub-double category of C generated by X. We say that a collection of 2-morphisms X of double category C generates C if C is equal to complete sub-double category of C generated by X. We will say that a double category C is **globularily generated** if C is generated by its collection of globular 2-morphisms. We write **gCat** for full subcategory of category **dCat** generated by collection of globularily generated double categories. In [7] it is proven that category **gCat** is reflective in category **dCat**. A left adjoint γ to inclusion of **gCat** in **dCat** is explicitly constructed. The image γC of a double category C under functor γ is equal to sub-double category of C generated by collection of globular 2-morphisms of C. The image γF of a double functor F, with double category C as domain category, under functor γ is equal to restriction of F to double category γC. Functor γ is referred to as the globularily generated piece functor and is interpreted as an analog of horizontalization functor H introduced above.

Decorated bicategories

We say that a pair C, B, formed by a category C and a bicategory B, is a decorated bicategory if collection of objects of category C equals collection of 0-cells of B. In that case we will call B the underlying bicategory of decorated bicategory C, B and C the decoration of C, B. We will usually denote decorated bicategories by single scripted letters, we will identify decorated bicategories with their underlying bicategories, and we will indicate decorations with asterisks, e.g. if letter $\mathcal{B}$ stands for a decorated bicategory, letter

$\mathcal{B}$ will also denote underlying bicategory of $\mathcal{B}$, and $\mathcal{B}^*$ will denote decoration of $\mathcal{B}$. This convention will be used in general statements and proofs, but it will not be implemented in certain particular cases. Given a decorated bicategory $\mathcal{B}$, in analogy with notation used in the theory of double categories we will call 0-, 1-, and 2-cells of underlying 2-category of $\mathcal{B}$, objects, horizontal morphisms, and globular morphisms of $\mathcal{B}$ respectively. We will call morphisms of decoration $\mathcal{B}^*$ of $\mathcal{B}$ vertical morphisms of $\mathcal{B}$. Given decorated bicategories $\mathcal{B}$ and $\mathcal{B}'$ we say that a pair F, F^* is a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$ if F is a bifunctor from underlying bicategory of $\mathcal{B}$ to underlying bicategory of $\mathcal{B}'$, F^* is a functor from decoration $\mathcal{B}^*$ of $\mathcal{B}$ to decoration $\mathcal{B}'^*$ of $\mathcal{B}$, and object functions of F and F^* coincide. In this case we call F the underlying bifunctor of pair F, F^* and we call F^* the decoration of F, F^*. We will usually identify decorated bifunctors with their underlying bifunctors and we use asterisks to indicate decorations e.g. when letter F denotes a decorated bifunctor, we also let letter F denote underlying bifunctor of F, and we let F^* denote decoration of F. Exceptions to this convention will apply in certain cases. Composition of decorated bifunctors is performed component-wise. This composition operation is associative and unital and thus provides pair formed by collection of decorated bicategories and collection of decorated bifunctors with the structure of a category. We denote this category by $\mathbf{bCat}^*$. We extend the construction of horizontalization functor H to a functor from category $\mathbf{dCat}$ to category $\mathbf{bCat}^*$. Given a double category C, we denote by H^*C pair formed by horizontalization, HC, of C, and object category C_0 of C. Thus defined, H^*C is a decorated bicategory. Given a double functor F between double categories C and D we denote by H^*F pair formed by horizontalization HF of F and object functor F_0 of F. Thus defined, pair H^*F associated to double functor F is a decorated bifunctor from decorated horizontalization H^*C of C to decorated horizontalization H^*D of D. Function associating to every double category C decorated horizontalization H^*C of C, together with function associating to every double functor F decorated horizontalization H^*F of F, is a functor from category $\mathbf{dCat}$ to category $\mathbf{bCat}^*$. We denote this functor by H^* and we call it the decorated horizontalization functor. We keep writing H^* for restriction of horizontalization functor H^* to category $\mathbf{gCat}$ of globularily generated double categories. Given a decorated bicategory $\mathcal{B}$ we say that a double category C is a **verticalization** of $\mathcal{B}$ if $\mathcal{B}$ is equal to decorated horizontalization H^*C of C.

Given a globularily generated double category C we write H_1 for the union of collection of globular 2-morphisms of C and collection of horizontal identities of vertical morphisms of C and we write V_1 for subcategory of category of morphisms C_1 of C, generated by collection H_1. Now, given a positive integer k strictly greater than 1, if we assume category V_{k-1} has been defined we associate to V_{k-1} a category V_k. First denote by H_k collection of all possible horizontal compositions of 2-morphisms in category V_{k-1}. We make, in that case, category V_k to be subcategory of category of morphisms C_1 of C, generated by collection H_k. We thus associate, recursively, to every double category C, a sequence of subcategories V_k of category of morphisms C_1 of C. We call, for every k, category V_k the k-th vertical category associated to double category C. We have used, in the above construction, for every k, an auxiliary collection of 2-morphisms H_k of C. Observe that for each k collection H_k both contains the horizontal identity of every vertical morphism in C and is closed under horizontal composition. If double category C is strict, then, for every k, collection H_k is collection of morphisms of a category whose collection of objects is collection of vertical morphisms of C. In that case we call category H_k the k-th horizontal category associated to double category C. By the way sequence of vertical categories V_k was constructed it is easily seen that for every k inclusions

$$\mathrm{Hom}V_k \subseteq H_{k+1} \subseteq \mathrm{Hom}V_{k+1}$$

hold. This implies that k-th vertical category V_k associated to double category C is a subcategory of $k+1$-th vertical category V_{k+1} associated to C for every k. Moreover, in the case in which double category C is strict, k-th horizontal category H_k associated to C is a subcategory of $k+1$-th horizontal category H_{k+1} associated to C for every k. The following lemma [7] says that sequence of vertical categories of a globularily generated double category forms a filtration of its category of morphisms.

Lemma 2.1. *Let C be a globularily generated double category. Morphism category C_1 of C is equal to the limit $\varinjlim V_k$ in category adjacent to **Cat** of sequence V_k of vertical categories associated to C.*

Given a strict double category C pair τC formed by collection of vertical morphisms of C and collection of 2-morphisms of C is a category. Composition operation in pair τC is horizontal composition in C. We call category τC associated to a strict double category C the transversal category associated to C.

Corollary 2.2. *Let C be a globularily generated double category. If C is a strict double category, then transversal category τC associated to C is equal to the limit $\varinjlim H_k$, in category adjacent to **Cat**, of sequence of horizontal categories associated to C.*

Given a double category C we call sequence V_k of vertical categories associated to C the vertical filtration of C. In the case in which double category C is strict we call sequence of horizontal categories H_k associated to C the horizontal filtration associated to C. When vertical or horizontal filtrations of different double categories are studied we indicate the double category of which corresponding filtrations is being considered with the double category as superscript, e.g. we write V_k^C for the k-th vertical category of a double category C. The definitions of vertical and horizontal filtrations associated of a double category admit functorial extensions. This follows from the following lemma [7].

Lemma 2.3. *Let C and C' be globularily generated double categories. Let $G : C \to C'$ be a double functor from C to C'. Let k be a positive integer. The image of collection H_k^C under morphism function of morphism functor G_1 of G is contained in collection $H_k^{C'}$. The image of k-th vertical category V_k^C associated to C under morphism functor G_1 of G is contained in k-th vertical category $V_k^{C'}$ associated to C'.*

Given a double functor $G : C \to C'$ from a double category C to a double category C' and k a positive integer, we write H_k^G for restriction to collection H_k^C of morphism function of morphism functor G_1 of G. We call H_k^G the k-th horizontal function associated to G. Assuming C and C' are strict double categories, we call pair formed by morphism function of object functor G_0 of G and k-th horizontal function H_k^G the k-th horizontal functor associated to G. Finally, we write V_k^G for restriction, to k-th vertical category V_k^C associated to C of morphism functor G_1 of G. Thus defined V_k^G is, by the above lemma, a functor from k-th vertical category V_k^C associated to C to k-th vertical category $V_k^{C'}$ associated to C'. We call V_k^G the k-th vertical functor associated to G.

Faithfulness of decorated horizontalization

We now prove that decorated horizontalization functor H^* is faithful when restricted to category **gCat** of globularily generated double categories. We begin with the following proposition.

Proposition 2.4. *Let C and C' be globularily generated double categories. Let $G, G' : C \to C'$ be double functors. Suppose decorated horizontalizations H^*G and H^*G' of G and G' are equal. In that case for every positive integer k, k-th horizontal functions H_k^G and $H_k^{G'}$ associated to G and G' respectively are equal, and k-th vertical functors V_k^G and $V_k^{G'}$ associated to G and G' respectively are equal.*

Proof. Let C and C' be globularily generated double categories. Let $G, G' : C \to C'$ be double functors. Suppose decorated horizontalizations H^*G and H^*G' of G and G' are equal. Let k be a positive integer. We wish to prove in that case that k-th horizontal functions H_k^G and $H_k^{G'}$ associated to G and G' respectively are equal, and that k-th vertical functors V_k^G and $V_k^{G'}$ associated to G and G' respectively are equal.

We proceed by induction on k. We prove first that first horizontal functions H_1^G and $H_1^{G'}$ associated to G and G' are equal. Let Φ be a globular morphism in C. The image $H_1^G\Phi$ of Φ under function H_1^G is equal to the image $G\Phi$ of Φ under G. This is equal, given that Φ is globular, to image $H^*G\Phi$ of Φ under decorated horizontalization H^*G of G, which is equal, by the assumption of the lemma, to the image $H^*G'\Phi$ of Φ under decorated horizontalization H^*G' of G', and this is equal to the image $H_1^{G'}\Phi$ of Φ under function $H_1^{G'}$. This, together the fact that both G and G' are double functors and thus preserve horizontal identities implies that functions H_1^G and $H_1^{G'}$ are equal. We now prove that functors V_1^G and $V_1^{G'}$ are equal. Observe first that restriction of morphism function of functor V_1^G to collection H_1^C is equal to function H_1^G and that restriction of morphism function of functor $V_1^{G'}$ to collection H_1^C is equal to function $H_1^{G'}$. This, the previous argument, the fact that collection H_1^C generates category V_1^C, and functoriality of V_1^G and $V_1^{G'}$, implies that functors V_1^G and $V_1^{G'}$ are equal.

Let now k be a positive integer strictly greater than 1. Suppose that for every positive integer n strictly less than k functions H_n^G and $H_n^{G'}$ are equal and functors V_n^G and $V_n^{G'}$ are equal. We prove now that under these assumptions functions H_k^G and $H_k^{G'}$ are equal. Observe first that restriction of function H_k^G to collection $\mathrm{Hom}_{V_{k-1}^C}$ of morphisms of category V_{k-1}^C is equal to morphism function of functor V_{k-1}^G and that restriction of function $H_k^{G'}$ to collection $\mathrm{Hom}_{V_{k-1}^C}$ of morphisms of V_{k-1}^C is equal to morphism function of functor $V_{k-1}^{G'}$. This, together with induction hypothesis, and the fact that G and G' intertwine source and target functors and horizontal composition bifunctors of C and C' implies that functions H_k^G and $H_k^{G'}$ are equal. We now prove that functors V_k^G and $V_k^{G'}$ are equal. Observe that restriction of

morphism function of functor V_k^G to collection H_k^C is equal to function H_k^G and that restriction of morphism function of functor $V_k^{G'}$ to collection H_k^C is equal to function $H_k^{G'}$. This, together with the previous argument, the fact that collection H_k^C generates category V_k^C, and functoriality of both V_k^G and $V_k^{G'}$ proves that functors V_k^G and $V_k^{G'}$ are equal. This concludes the proof. ■

Corollary 2.5. *Let C and C' be double categories. Suppose C is globularily generated. Let $G, G' : C \to C'$ be double functors. If decorated horizontalizations H^*G and H^*G' associated to G and G' respectively are equal, then double functors G and G' are equal.*

Proof. Let C and C' be double categories. Let $G, G' : C \to C'$ be double functors. Suppose that C is globularily generated and that decorated horizontalizations H^*G and H^*G' of G and G' are equal. We wish to prove, in this case, that double functors G and G' are equal.

Globular pieces γG and $\gamma G'$ of G and G' are both double functors from globularily generated double category C to globularily generated piece $\gamma C'$ of double category C'. Decorated horizontalization $H^*\gamma G$ of γG is equal to decorated horizontalization H^*G of G and decorated horizontalization $H^*\gamma G'$ of $\gamma G'$ is equal to decorated horizontalization H^*G' of G'. It follows, from the assumption of the corollary that double functors γG and $\gamma G'$ satisfy the assumptions of proposition ? and thus for every positive integer k k-th term $V_k^{\gamma G}$ of vertical filtration associated to γG and k-th term $V_k^{\gamma G'}$ of vertical filtration associated to $\gamma G'$ are equal. By ? both γG and $\gamma G'$ admit decompositions as limits $\varinjlim V_k^{\gamma G}$ and $\varinjlim V_k^{\gamma G'}$ of vertical filtrations $V_k^{\gamma G}$ and $V_k^{\gamma G'}$ associated to G and G' respectively. It follows, from this, that globular piece γG of G and globular piece $\gamma G'$ of G' are equal. Finally, globular piece double functors γG and $\gamma G'$ associated to G and G' are equal to codomain restrictions, to globular piece dobule category $\gamma C'$ of C' by the assumption that C is globularily generated and thus double functors γG and $\gamma G'$ are equal if and only if double functors G and G' are equal. This concludes the proof. ■

Corollary 2.6. *Restriction to category **gCat** of decorated horizontalization functor H^* is faithful.*

Examples

We now present examples of double categories and decorated bicategories. The following will be the only examples used throughout the paper.

Trivially decorated bicategories: Let B be a bicategory. We write B^* for discrete category generated by collection of 0-cells B_0 of B. Pair B^*, B is a decorated bicategory. We call this decorated bicategory the trivially decorated bicategory associated to bicategory B and we keep denoting it by B. Observe that decorated horizontalization $H^*\overline{B}$ of trivial double category $\overline{B}$ associated to trivially decorated bicategory associated to B is equal to trivially decorated bicategory B, that is, trivial double category $\overline{B}$ associated to trivially decorated bicategory B is a verticalization of B.

Single object decorated bicategories: Let C be a monoidal category. We denote by $2C$ bicategory with a single 0-cell a and whose collections of 1- and 2-cells are collection of objects of C, and collection of morphisms of C respectively. Horizontal composition in $2C$ is defined by tensor product of C, vertical composition in $2C$ is defined by composition of morphisms of C, and identity 1-cell of 0-cell a is monoidal unit 1 of C. Exchange relation in $2C$ follows from functoriality of monoidal structure of C. Left and right identity transformations and associator in $2C$ are defined by left and right unit transformations and associator of C. Bicategory $2C$ is a 2-category if and only if monoidal category C is strict. Let A be a monoid. We denote by mA category with single object a, with monoid of endomorphisms in mA equal to A. We denote by $2C_A$ pair $mA, 2C$, formed by category mA associated to A, and by 2-category $2C$ associated to C. $2C_A$ thus defined is a decorated bicategory. Every single object decorated bicategory is of this form.

The following two examples illustrate the action of decorated horizontalization functor. They will not be used in the rest of the paper.

Algebras: Let $\mathbf{Alg}_0$ denote category whose collection of objects is collection of complex algebras and whose collection of morphisms is collection of unital algebra morphisms. Given algebras A, B, C and D, left-right A-B bimodule M and left-right C-D-bimodule N, we say that a triple (f, Φ, g), where f is a unital algebra morphism from A to C, g is a unital algebra morphism from B to D and Φ is a linear transformation from M to N; is an equivariant bimodule morphism from M to N, if for every $a \in A$, $b \in B$, and $x \in M$ equation

$\Phi(axb) = f(a)\Phi(x)g(b)$ holds. Composition of equivariant bimodule morphisms is performed entry-wise. Let $\mathbf{Alg}_1$ denote category whose collection of objects is collection of bimodules over complex algebras and whose collection of morphisms is collection of equivariant bimodule morphisms. Denote by $\mathbf{Alg}$ pair formed by categories $\mathbf{Alg}_0$ and $\mathbf{Alg}_1$. Denote by i functor from $\mathbf{Alg}_0$ to $\mathbf{Alg}_1$, associating algebra A as a left-right A-bimodule to every algebra A, and associating equivariant bimodule morphism (f, f, f) to every unital algebra morphism f. Given algebras A, B and C, left A-B bimodule M and left-right B-C bimodule N write $N * M$ for tensor product $M \otimes_B N$ relative to B. Given equivariant bimodule morphisms (f, Φ, g) and (g, Ψ, h) we write $(g, \Psi, h) * (f, \Phi, g)$ for tensor product $(f, \Phi \otimes_g \Psi, h)$. Denote by $*$ bifunctor, from $\mathbf{Alg}_1 \times_{\mathbf{Alg}_0} \mathbf{Alg}_1$ to $\mathbf{Alg}_1$ defined by operations $*$ defined above. This structure provides pair $\mathbf{Alg}$ of categories $\mathbf{Alg}_0$ and $\mathbf{Alg}_1$ with the structure of a double category. Left and right identity transformations and associator in $\mathbf{Alg}$ are defined by those of relative tensor product bifunctor. Decorated horizontalization $H^*\mathbf{Alg}$ is equal to bicategory $\underline{\mathbf{Alg}}$ of unital algebras, algebra bimodules, and bimodules morphisms presented in [9], decorated by category $\mathbf{Alg}_0$ of unital algebra morphisms.

Cobordisms: Let n be a positive integer. Let $\mathbf{Cob}(n)_0$ denote category whose collection of objects is collection of closed n-dimensional smooth manifolds and whose collection of morphisms is collection of diffeomorphisms between manifolds. Given closed n-manifolds X, Y, Z and W, a cobordism M from X to Y, and a cobordism N from Z to W, we will say that a triple (f, Φ, g), where f is a diffeomorphisms from X to Z, g is a diffeomorphism from Y to W, and where Φ is a diffeomorphism from M to N; is an equivariant diffeomorphism from M to N if restriction of Φ to X equals f and restriction of Φ to Y equals g. Composition of equivariant morphisms between cobordisms is performed entry-wise. Let $\mathbf{Cob}(n)_1$ denote category whose collection of objects is collection of cobordisms between closed n-dimensional manifolds and whose collection of morphisms is collection of equivariant diffeomorphisms between cobordisms. Given an n-dimensional manifold X we write i_X for cobordism $X \times [0, 1]$. Given a diffeomorphism $f : X \to Y$ between closed n-dimensional manifolds X and Y, we denote by i_f equivariant diffeomorphism $(f, f \times [0, 1], f)$ from cobordism i_X to cobordism i_Y. These two functions define a functor from $\mathbf{Cob}(n)_0$ to $\mathbf{Cob}(n)_1$. Denote this functor by i. Given compatible cobordisms M and N with respect to manifold X we write $N * M$ for joint union $M \cup_X Y$ with respect to X. Finally, given equivariant diffeomorphisms (f, Φ, g) and (g, Ψ, h), com-

patible with respect to diffeomorphism g denote by $(g, \Psi, h) * (f, \Phi, g)$ joint union $(f, \Phi \cup_g \Psi, h)$ of (f, Φ, g) and (g, Ψ, h) with respect to g. The two operations defined above form a bifunctor which we denote by $*$. With this structure pair $\mathbf{Cob}(n)$ is a double category, where identity transformations and associator come from the obvious diffeomorphisms from cobordism theory. Decorated horizontalization $H^*\mathbf{Cob}(n)$ of $\mathbf{Cob}(n)$ is equal to bicategory $\underline{\mathbf{Cob}(n)}$ of n-dimensional cobordisms and diffeomorphisms presented in [9] decorated by grupoid of diffeomorphisms of closed dimensional n-manifolds.

Outlook

In [1] the problem of extending function associating, to every von Neumann algebra A its Haagerup standard form construction L^2A to a functor from category of von Neumann algebras and von Neumann algebra morphisms was proposed. Moreover, it was asked if function associating, to every horizontally composable pair of bimodules ${}_AH_B$ and ${}_BK_C$ over von Neumann algebras A, B, and C, Connes' fusion $H \boxtimes_B K$ of H and K along B, is functorial with respect to algebra B. Both these problems were positively solved in [1] for the case in which von Neumann algebras and von Neumann algebra morphisms considered had finite dimensional center and were finite respectively. These results can be phrased as follows: Denote by $\mathbf{vN}^f$ 2-category of von Neumann algebras with finite dimensional center, von Neumann algebra bimodules, and intertwiners, decorated by category $\mathbf{vN}_0^f$, whose collection of objects is collection of von Neumann algebras with finite dimensional center, and whose collection of morphisms is collection of finite von Neumann algebra morphisms. Horizontal identity in 2-category $\mathbf{vN}^f$ is given by Haagerup's L^2-space construction and horizontal composition is given by Connes' fusion operation. Results in [1] are equivalent to the fact that there exists a verticalization of decorated 2-category $\mathbf{vN}^f$, analogous to double category **Alg** of unital algebras and bimodules, having collection of equivariant von Neumann algebra morphisms as collection of 2-morphisms. Problem of general categorification of Haagerup's L^2-space construction and Connes' fusion operation is equivalent to the problem of finding a verticalization of 2-category of general von Neumann algebras, von Neumann algebra bimodules, and intertwiners, decorated by category of von Neumann algebras and general von Neumann algebra morphisms, and such that its collection of 2-morphisms is precisely collection of equivariant bimodule morphisms. The author considers the study of globularily generated verticalizations of decorated 2-categories as a preamble to problems similar to problem described above.

3 Free globularily generated double category

In this section we introduce our main object of study. Namely, we introduce the free globularily generated double category construction. We do this in several steps. We begin with a few preliminary definitions and results.

Evaluations

Given collections X and Y, functions $s, t : X \to Y$ from X to Y and a sequence $x_1, ..., x_n$ of elements of X, we will say that sequence $x_1, ..., x_k$ is compatible with respect to pair of functions s, t if equation $tx_{i+1} = sx_i$ is satisfied for every positive integer i less than or equal to $k-1$. Equivalently, sequence $x_1, ..., x_k$ is compatible with respect to pair s, t if sequence $x_1, ..., x_n$ is a composable sequence of morphisms in free category generated by X with s and t as domain and codomain functions respectively. In this case, given a compatible sequence $x_1, ..., x_k$ of elements of collection X, with respect to pair of functions s, t, we call any way of writing word $x_k...x_1$ following an admissible parenthesis pattern, an evaluation of sequence $x_1, ..., x_k$. Equivalently, evaluations of a compatible sequence $x_1, ..., x_k$ are different ways of writing word $x_k...x_1$ composing elements of sequence $x_1, ..., x_k$ two by two in free category generated by X, with functions s and t as domain and codomain functions. For example, (yx) is the only evaluation of a 2 term compatible sequence x, y and $(x(yz)), ((xy)z)$ are the two evaluations of a compatible 3 term compatible sequence x, y, z. We will write $X_{s,t}$ for collection of evaluations of finite sequences of elements of class X, compatible with respect to pair of functions s, t.

Given functions $s, t : X \to Y$ from a collection X to a collection Y, we write $\tilde{s}$ and $\tilde{t}$ for functions $\tilde{s}, \tilde{t} : X_{s,t} \to Y$ defined as follows: Given an evaluation Φ of a sequence $x_1, ..., x_k$ of elements of X, compatible with respect to s and t, we make $\tilde{s}\Phi$ and $\tilde{t}\Phi$ to be equal to sx_1 and tx_k respectively. Observe that $\tilde{s}\Phi$ and $\tilde{t}\Phi$ is equal for every evaluation of compatible sequence $x_1, ..., x_k$. Given a pair of sequences $x_1, ..., x_k$ and $x_{k+1}, ..., x_n$ of elements of X, compatible with respect to s and t, such that 2 term sequence x_k, x_{k+1} is compatible with respect to s and t, and given evaluations Φ and Ψ of sequences $x_1, ..., x_k$ and $x_{k+1}, ..., x_n$ respectively, equation $\tilde{t}\Phi = \tilde{s}\Psi$ is satisfied and concatenation of Ψ and Φ defines an evaluation of compatible sequence $x_1, ..., x_k, x_{k+1}, ..., x_n$. We denote concatenation of evaluations Φ and Ψ satisfying conditions above by $\Psi *_{s,t} \Phi$. These operations define a function from $X_{s,t} \times_Y X_{s,t}$ to $X_{s,t}$ where fibration in $X_{s,t} \times_Y X_{s,t}$ is taken with respect to pair $\tilde{s}, \tilde{t}$. We write $*_{s,t}$ for this function.

Now, given collections X, X', Y, and Y', and functions $s, t : X \to Y$ from X to Y and $s', t' : X' \to Y'$ from X' to Y', we say that a pair of functions $\varphi : X \to X'$ and $\phi : Y \to Y'$ is compatible if the following two squares commute



Given collections X, X', Y, and Y', functions $s, t : X \to Y$ and $s', t' : X' \to Y'$, and a compatible pair of functions $\varphi : X \to X'$ and $\phi : Y \to Y'$, if $x_1, ..., x_k$ is a sequence of elements of X, compatible with respect to pair of functions s, t, then in that case sequence $\varphi x_1, ..., \varphi x_k$ is compatible with respect to s' and t'. Moreover, given an evaluation Φ of compatible sequence $x_1, ..., x_k$, the same parenthesis pattern defining evaluation Φ defines an evaluation of compatible sequence $\varphi x_1, ..., \varphi x_k$. We write $\mu_{\varphi,\phi}\Phi$ for this evaluation. We write $\mu_{\varphi,\phi}$ for function from collection $X_{s,t}$ of evaluations with respect to triple $X_{s,t}$ to collection $X'_{s',t'}$ of evaluations with respect to triple X', s', t' associating evaluation $\mu_{\varphi,\phi}\Phi$ to every evaluation Φ in $X_{s,t}$. The proof of the following lemma is straightforward.

Lemma 3.1. *Let X, X', Y, and Y' be classes. Let $s, t : X \to Y$ and $s', t' : X' \to Y'$ be functions. Let $\varphi : X \to X'$ and $\phi : Y \to Y'$ be functions such that pair φ, ϕ is compatible. In that case function $\mu_{\varphi,\phi}$ associated to pair φ, ϕ satisfies the following conditions*

1. *The following two squares commute*



2. *The following square commutes*



Notational conventions

Given a vertical morphism α in a decorated bicategory $\mathcal{B}$, we will write i_α for singleton $\{\alpha\}$. We call i_α the formal horizontal identity of vertical morphism α. We will write $\mathbb{G}$ for the union of collection of globular morphisms in $\mathcal{B}$ and collection of formal horizontal identities i_α with α running over collection of vertical morphisms of $\mathcal{B}$. We will further adopt the following notational conventions.

1. Let Φ be a globular morphism in $\mathcal{B}$. We denote by $d_0\Phi$ and $c_0\Phi$ domain and codomain of Φ in $\mathcal{B}$ respectively. Let α be a vertical morphism in $\mathcal{B}$. Let a and b be domain and codomain, in decoration $\mathcal{B}^*$ of α. In that case we write $d_0 i_\alpha$ and $c_0 i_\alpha$ for horizontal identities id_a and id_b of a and b in $\mathcal{B}$ respectively. We write d_0 and c_0 for functions from collection $\mathbb{G}$ to collection $\mathcal{B}_1$ of horizontal morphisms of $\mathcal{B}$ associating $d_0\Phi$ and $c_0\Phi$ to every element Φ of $\mathbb{G}$.

2. Let Φ be a globular morphism in $\mathcal{B}$. We denote by $s_0\Phi$ and $t_0\Phi$ source and target of Φ in $\mathcal{B}$. We extend the definition of source and target of globular morphisms in $\mathcal{B}$ to formal horizontal identities of vertical morphisms in $\mathcal{B}$ as follows. Let α be a vertical morphism in $\mathcal{B}$. In that case we make both $s_0 i_\alpha$ and $t_0 i_\alpha$ to be equal to α. We write s_0 and t_0 for functions from $\mathbb{G}$ to collection of vertical morphisms of $\mathcal{B}$ associating $s_0\Phi$ and $t_0\Phi$ to every element Φ of $\mathbb{G}$.

Functions d_0, c_0, s_0, and t_0 defined above are easily seen to be related by the following conditions

1. The following two triangles commute



2. The following two squares commute



Given a decorated bicategory $\mathcal{B}$, we denote by p function from collection of evaluations $\mathcal{B}_{1_{d,c}}$ of collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$, with respect to pair formed by domain and codomain functions d, c in adjacent bicategory $\mathcal{B}$, to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$ such that for every composable sequence $f_1, ..., f_k$ of horizontal morphisms in $\mathcal{B}$ and for every evaluation Φ of $f_1, ..., f_k$, the image $p\Phi$ of evaluation Φ under p is equal to horizontal composition of sequence $f_1, ..., f_k$ in $\mathcal{B}$ following parenthesis pattern defining Φ. We call p the projection associated to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$. We now proceed to the construction of the main object of study of this section.

Main construction

Given a decorated bicategory $\mathcal{B}$ we write E_1 for collection of evaluations $\mathbb{G}_{s_0,t_0}$ of collection $\mathbb{G}$ defined above with respect to pair of functions s_0, t_0. We denote by s_1 and t_1 functions $\tilde{s}_0$ and $\tilde{t}_0$ defined by s_0 and t_0. Thus defined s_1 and t_1 are functions from E_1 to collection $\mathrm{Hom}_{\mathcal{B}^*}$ of vertical morphisms of $\mathcal{B}$. We write $*_1$ for operation $*_{s_1,t_1}$ on E_1. Finally, we write d_1 for composition $p\mu_{d_0,id}$ of function $\mu_{d_0,id}$ associated to pair d_0, id and projection p associated to $\mathcal{B}_1$, and we write c_1 for composition $p\mu_{c_0,id}$ of function $\mu_{c_0,id}$ associated to pair c_0, id and projection p from associated to $\mathcal{B}_1$. Thus defined d_1 and c_1 are functions from E_1 to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$.

The following theorem is the first step towards the free globularily generated double category construction.

Theorem 3.2. *Let $\mathcal{B}$ be a decorated bicategory. There exists a pair of sequences of triples E_k, d_k, c_k and F_k, s_{k+1}, t_{k+1} with k running through collection of positive integers, such that for each positive integer k, E_k is a collection containing collection E_1 defined above, d_k, c_k are functions from E_k to collection $\mathcal{B}_1$ of horizontal morphisms of $\mathcal{B}$ extending functions d_1 and c_1 defined above respectively, F_k is a category having collection $\mathcal{B}_1$ of horizontal morphisms of $\mathcal{B}$ as collection of objects, and s_{k+1}, t_{k+1} are functors from F_k to decoration $\mathcal{B}^*$ of $\mathcal{B}$. Moreover, pair of sequences E_k, d_k, c_k and F_k, s_{k+1}, t_{k+1} satisfies the following conditions*

1. *For every positive integer k collection of morphisms Hom_{F_k} of category F_k is contained in collection E_{k+1}. Moreover, collection E_{k+1} is equal to collection of evaluations $Hom_{F_k s_{k+1}, t_{k+1}}$ of collection Hom_{F_k} of morphisms of F_k with respect to pair formed by morphism functions of functors s_{k+1}, t_{k+1}.*

2. *For every positive integer k collection E_k is contained in collection of morphisms Hom_{F_k} of category F_k. Moreover, category F_k is equal to free category generated by collection E_k with functions d_k and c_k as domain and codomain functions respectively. Restriction of morphism functions of functors s_{k+1} and t_{k+1} to collection E_1 is equal to functions s_1 and t_1 defined above.*

3. *For every positive integer k the following four triangles commute*



Conditions 1-3 above determine pair of sequences of triples E_k, d_k, c_k and F_k, s_{k+1}, t_{k+1}.

Proof. Let $\mathcal{B}$ be a decorated bicategory. We wish to construct a pair of sequences of triples E_k, d_k, c_k and F_k, s_{k+1}, t_{k+1} with k running through collection of positive integers, such that for each positive integer k, E_k is a

collection extending collection E_1 associated to $\mathcal{B}$, such that d_k and c_k are functions from E_k to collection $\mathcal{B}_1$ of horizontal morphisms of $\mathcal{B}$ extending functions d_1 and c_1, such that F_k is a category having collection $\mathcal{B}_1$ of horizontal morphisms of $\mathcal{B}$ as collection of objects and s_{k+1}, t_{k+1} are functors from F_k to decoration $\mathcal{B}^*$ of $\mathcal{B}$. Moreover, we wish to define pair of sequences E_k, d_k, c_k and F_k, s_{k+1}, t_{k+1} in such a way that conditions 1-3 above are satisfied.

We proceed by induction on k. We begin by defining triple F_1, s_2, t_2. We make F_1 to be free category generated by collection E_1 with respect to functions d_1, c_1. Functions s_1 and t_1 are compatible with domain and codomain functions d_1 and c_1 and thus admit a unique extension to functors from category F_1 to decoration $\mathcal{B}^*$ of $\mathcal{B}$. We make s_2 and t_2 corresponding functorial extensions of s_1 and t_1 respectively. We write $\bullet_1$ for composition operation in F_1.

Let now k be a positive integer strictly greater than 1. Assume we have extended the definition of triple E_1, d_1, c_1 and the definition of triple F_1, s_2, t_2 of the previous paragraph, to a sequence of pairs of triples E_m, d_m, c_m and F_m, s_{m+1}, t_{m+1} with m running through collection of positive integers strictly less than k, where $E_m^{\mathcal{B}}$ is assumed to be a collection containing E_1, d_m and c_m are assumed to be functions from E_m to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$ extending functions d_1 and c_1 respectively, F_m is assumed to be a category containing F_1 as subcategory, and s_{m+1} and t_{m+1} are assumed to be functors from F_m to decoration $\mathcal{B}^*$ of $\mathcal{B}$ extending functors s_2 and t_2 respectively. Moreover, we assume that pair of sequences of triples E_m, d_m, c_m and F_m, s_{m+1}, t_{m+1} satisfy conditions 1-3 above.

We now wish to extend the definition of pair of sequences of triples $E_m^{\mathcal{B}}, d_m, c_m$ and $F_m^{\mathcal{B}}, s_{m+1}, t_{m+1}$ to the definition of pair of triples $E_k^{\mathcal{B}}, d_k, c_k$ and $F_k^{\mathcal{B}}, s_{k+1}, t_{k+1}$ satisfying the conditions of the theorem. We begin with the definition of triple E_k, d_k, c_k. We make E_k to be collection of evaluations $\mathrm{Hom}_{F_{k-1}{}_{s_k,t_k}}$ of collection of morphisms $\mathrm{Hom}_{F_{k-1}}$ of category F_{k-1} with respect to pair formed by morphism functions of functors s_k and t_k. We write s_{k+1} and t_{k+1} for extensions $\tilde{s}_k$ and $\tilde{t}_k$, to collection E_k, of morphism functions of s_k and t_k. We denote by $*_k$ concatenation operation $*_{s_{k+1},t_{k+1}}$ in E_k, with respect to extensions s_{k+1} and t_{k+1}. We now make function d_k to be composition $p\mu_{d_{k-1},id}$ of function associated to pair formed by domain function d_{k-1} in F_{k-1} and identity function in collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$ and projection p associated to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$. We make function c_k to be composition $p\mu_{c_{k-1},id}$ of function associated to pair formed by codomain function in F_{k-1} and identity function in

collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$, and projection p associated to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$. Functions d_k and c_k are well definied. Thus defined d_k and c_k are functions from collection E_k to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$ satisfying condition 3 of the theorem by lemma?. We now define triple F_k, s_{k+1}, t_{k+1}. We make category F_k to be free category generated by collection E_k defined above, with functions d_k and c_k as domain and codomain functions respectively. Collection of objects of category F_k is thus collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$. We write $\bullet_k$ for composition operation in F_k. By the fact that functions d_k, c_k, s_k and t_k satisfy condition 3 of the theorem it follows that pairs formed by functions s_k and t_k together with domain and codomain functions defined on collection of horizontal morphisms of $\mathcal{B}$ admit unique extensions to functors from category F_k to decoration $\mathcal{B}_1$ of $\mathcal{B}$. We make s_{k+1} and t_{k+1} to be these functors. Thus defined triple F_k, s_{k+1}, t_{k+1} satisfies the conditions of the theorem.

It is obvious that conditions 1-3 of the theorem determine pair of sequences E_k, d_k, c_k and F_k, s_{k+1}, t_{k+1}. This concludes the proof. ■

Observation 3.3. Let $\mathcal{B}$ be a decorated bicategory. Let m, k be positive integers such that m is less than or equal to k. In that case, as defined above, collection E_m is contained in collection E_k, functions d_m and c_m are equal to restrictions to collection E_k of functions d_k and c_k respectively, and concatenation operation $*_m$ is equal to restriction to E_m of operation $*_k$. Moreover, category F_m is a subcategory of category F_k and functors s_{m+1} and t_{m+1} are equal to restrictions to category F_m of functors s_{k+1} and t_{k+1} respectively.

Observation 3.4. Let $\mathcal{B}$ be a decorated bicategory. Let k be a positive integer. The following two squares commute where q_1^k and q_2^k denote the left and right projections of $E_k \times_{\mathrm{Hom}_{\mathcal{B}^*}} E_k$ onto E_k respectively.

$$\begin{array}{ccc} E_k \times_{\mathrm{Hom}_{\mathcal{B}^*}} E_k & \xrightarrow{*_k} & E_k \\ \downarrow{\scriptstyle q_1^k} & & \downarrow{\scriptstyle s_k} \\ E_k & \xrightarrow[s_k]{} & \mathrm{Hom}_{\mathcal{B}^*} \end{array} \qquad \begin{array}{ccc} E_k \times_{\mathrm{Hom}_{\mathcal{B}^*}} E_k & \xrightarrow{*_k} & E_k \\ \downarrow{\scriptstyle q_2^k} & & \downarrow{\scriptstyle t_k} \\ E_k & \xrightarrow[s_k]{} & \mathrm{Hom}_{\mathcal{B}^*} \end{array}$$

Definition 3.5. Let $\mathcal{B}$ be a decorated bicategory. In that case we write E_∞ for the union $\bigcup_{k=1}^{\infty} E_k$ of sequence of collections E_k associated to $\mathcal{B}$. We

write d_∞ and c_∞ for the limits in category **Set** of sequences of functions d_k and c_k respectively. Thus defined d_∞ and c_∞ are functions from collection E_∞ to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$. We denote by $*_\infty$ the limit $\varinjlim *_k$ in category **Set** of sequence of operations $*_k$ defined in theorem ?. Thus defined $*_\infty$ is a function from $E_\infty \times_{\mathcal{B}^*} E_\infty$ to E_∞, where fibration in $E_\infty \times_{\mathcal{B}^*} E_\infty$ is taken with respect to pair of functions s_∞, t_∞. We write F_∞ for the limit $\varinjlim F_k$ in 2-category **Cat** of sequence of categories F_k. Collection of objects of category F_∞ is thus collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$, and collection of morphisms of F_∞ is equal to collection E_∞. Domain and codomain functions of category F_∞ are equal to functions d_∞ and c_∞. We write s_∞, t_∞ for the limits of sequences of functors s_k, t_k. Thus defined s_∞ and t_∞ are functors from category F_∞ to decoration $\mathcal{B}^*$ of $\mathcal{B}$. Finally, we write $\bullet_\infty$ for composition operation of F_∞. Thus defined $\bullet_\infty$ is the limit $\varinjlim \bullet_k$, in category **Set**, of sequence of composition operations $\bullet_k$ defining categories F_k.

Definition 3.6. Let $\mathcal{B}$ be a decorated bicategory. We write R_∞ for equivalence relation generated by the following relations defined on collection of morphisms E_∞ of category F_∞ associated to $\mathcal{B}$.

1. Let Φ_i, Ψ_i, $i = 1, 2$ be morphisms in F_∞ such that pairs Φ_1, Φ_2 and Ψ_1, Ψ_2 are compatible with respect to pair s_∞, t_∞ and such that pairs Φ_i, Ψ_i, $i = 1, 2$ are both compatible with respect to pair d_∞, c_∞. In that case we identify compositions:

$$(\Psi_2 *_\infty \Psi_1) \bullet_\infty (\Phi_2 *_\infty \Phi_1) \text{ and } (\Psi_2 \bullet_\infty \Phi_2) *_\infty (\Psi_1 \bullet_\infty \Phi_1)$$

2. Let Φ and Ψ be globular 2-morphisms of $\mathcal{B}$ such that pair Φ, Ψ is compatible with respect to pair s_∞, t_∞. In that case we identify composition $\Psi \bullet_\infty \Phi$ of Φ and Ψ with vertical compostion $\Psi\Phi$ of Φ and Ψ in $\mathcal{B}$. Moreover, if α and β are vertical morphisms in $\mathcal{B}$ such that pair α, β is composable in decoration $\mathcal{B}^*$ of $\mathcal{B}$, then, in that case, we identify composition $i_\beta \bullet_\infty i_\alpha$ of horizontal identities i_α and i_β with horizontal identity $i_{\beta\alpha}$ of composition $\beta\alpha$.

3. Let Φ and Ψ be globular 2-morphisms of $\mathcal{B}$ such that pair Φ, Ψ is compatible with respect to functors s_∞, t_∞. In that case we identify composition $\Psi *_\infty \Phi$ with horizontal composition $\Psi\Phi$ of Φ and Ψ.

4. Let Φ be a morphism in F_∞. We identify Φ with compositions

$$\lambda_{c_\infty\Phi} \bullet_\infty (\Phi *_\infty i_{t_\infty\Phi}) \bullet_\infty \lambda^{-1}_{d_\infty\Phi} \text{ and } \rho_{c_\infty\Phi} \bullet_\infty (i_{s_\infty\Phi} *_\infty \Phi) \bullet_\infty \rho^{-1}_{d_\infty\Phi}$$

where λ and ρ denote left and right identity transformations of bicategory adjacent to $\mathcal{B}$.

5. Let Φ, Ψ, Θ be elements of E_∞ such that triple Φ, Ψ, Θ is compatible with respect to pair s_∞, t_∞. In that case we identitfy compositions

$$A_{c_\infty\Phi, c_\infty\Psi, c_\infty\Theta} \bullet_\infty [\Theta *_\infty (\Psi *_\infty \Phi)] \text{ and } [(\Theta *_\infty \Psi) *_\infty \Phi] \bullet_\infty A_{d_\infty\Phi, d_\infty\Psi, d_\infty\Theta}$$

where A denotes the associator of bicategory adjacent to $\mathcal{B}$.

Lemma 3.7. *Let $\mathcal{B}$ be a decorated bicategory. In that case equivalence relation R_∞ is compatible with domain, codomain, and composition operation functions d_∞, c_∞, and $\bullet_\infty$ defining category structure on category F_∞ associated to $\mathcal{B}$.*

Proof. Let $\mathcal{B}$ be a decorated bicategory. We wish to prove that equivalence relation R_∞ is compatible with domain, codomain, and composition operation functions d_∞, c_∞, and $\bullet_\infty$ defining category structure on F_∞.

Let Φ_i, Ψ_i, $i = 1, 2$ be morphisms in F_∞ such that pairs Φ_1, Φ_2 and Ψ_1, Ψ_2 are compatible with respect to source and target functors s_∞, t_∞ of F_∞ and such that pairs Φ_i, Ψ_i, $i = 1, 2$ are compatible with respect to domain and codomain functions d_∞, c_∞. In that case domain

$$d_\infty(\Psi_1 *_\infty \Psi_1) \bullet_\infty (\Phi_2 *_\infty \Phi_1)$$

of morphism $(\Psi_1 *_\infty \Psi_1) \bullet_\infty (\Phi_2 *_\infty \Phi_1)$ is equal to domain $d_\infty(\Phi_2 *_\infty \Phi_1)$ of $(\Phi_2 *_\infty \Phi_1)$ which in turn is equal to composition $d_\infty\Phi_2 d_\infty\Phi_1$ of domains $d_\infty\Phi_1$ and $d_\infty\Phi_2$ of Φ_1 and Ψ_1. Now, domain

$$d_\infty(\Psi_2 \bullet_\infty \Phi_2) *_\infty (\Psi_1 \bullet_\infty \Phi_1)$$

of composition $(\Psi_1 \bullet_\infty \Phi_1) *_\infty (\Psi_2 \bullet_\infty \Phi_2)$ is equal to composition

$$d_\infty(\Psi_2 \bullet_\infty \Phi_2) d_\infty(\Psi_1 \bullet_\infty \Phi_1)$$

of domain $d_\infty(\Psi_1 \bullet_\infty \Phi_1)$ of composition $\Psi_1 \bullet_\infty \Phi_1$ and domain $d_\infty(\Psi_2 \bullet_\infty \Phi_2)$ of composition $\Psi_2 \bullet_\infty \Phi_2$. Domain $d_\infty(\Psi_1 \bullet_\infty \Phi_1)$ of composition $\Psi_1 \bullet_\infty \Phi_1$

is equal to domain $d_\infty\Phi_1$ of Φ_1 and domain $d_\infty(\Psi_2 \bullet_\infty \Phi_2)$ of composition $\Psi_2 \bullet_\infty \Phi_2$ is equal to domain $d_\infty\Phi_2$ of Φ_2. Thus domain

$$d_\infty(\Psi_2 \bullet_\infty \Phi_2) *_\infty (\Psi_1 \bullet_\infty \Phi_1)$$

of composition $(\Psi_2 \bullet_\infty \Phi_2) *_\infty (\Psi_1 \bullet_\infty \Phi_1)$ is equal to composition $d_\infty d_\infty \Phi_2 d_\infty \Phi_1$ of domain $d_\infty\Phi_1$ of Φ_1 and domain $d_\infty\Phi_2$ of Φ_2. We conclude that equivalence relation 1 in the definition of relation R_∞ is compatible with respect to domain function d_∞ of category F_∞. A similar computation proves that relation 1 in the definition of R_∞ is compatible with respect to codomain function c_∞ of F_∞.

Let now Φ, Ψ be elements of $\mathbb{G}$. suppose pair Φ, Ψ is compatible with respect to domain and codomain functions d_∞ and c_∞ of F_∞. In that case domain and codomain $d_\infty\Psi \bullet_\infty \Phi$ and $\Psi \bullet_\infty \Phi$ of composition $\Psi \bullet_\infty \Phi$ are equal to domain $d_\infty\Phi$ of Φ and codomain $c_\infty\Psi$ of Ψ respectively. Domain and codomain $d_\infty\Psi\Phi$ and $c_\infty\Psi\Phi$ of vertical composition $\Psi\Phi$ of Φ and Ψ is equal to domain $d_\infty\Phi$ of Φ and codomain $c_\infty\Psi$ of Ψ. We conclude that relation 2 in the definition of equivalence relation R_∞ is compatible with domain and codomain functions d_∞ and c_∞ of category F_∞.

Let Φ and Ψ be globular morphisms in $\mathcal{B}$. Suppose pair Φ, Ψ is compatible with respect to morphism functions of functors s_∞ and t_∞. In that case domain $d_\infty\Psi *_\infty \Phi$ and codomain $c_\infty\Psi *_\infty \Phi$ of horizontal composition $\Psi *_\infty \Phi$ of Φ and Ψ are equal to compositions, in $\mathcal{B}$, $d_\infty\Psi d_\infty\Phi$ and $c_\infty\Psi d_\infty\Phi$ respectively. Now, domain $d_\infty\Psi * \Phi$ and codomain $c_\infty\Psi * \Phi$ of horizontal composition, in $\mathcal{B}$, of Φ and Ψ is equal to compositions $d_\infty\Phi d_\infty\Psi$ and $c_\infty\Phi c_\infty\Psi$ respectively. This proves that relation 3 in the definition of equivalence relation R_∞ is compatible with domain and codomain functions d_∞ and c_∞ of category F_∞.

Let now Φ be a general morphism in F_∞. In that case domain

$$d_\infty\lambda_{c_\infty\Phi} \bullet_\infty (\Phi *_\infty i_{t_\infty\Phi}) \bullet_\infty \lambda^{-1}_{d_\infty\Phi}$$

of composition $\lambda_{c_\infty\Phi} \bullet_\infty (\Phi *_\infty i_{t_\infty\Phi}) \bullet_\infty \lambda^{-1}_{d_\infty\Phi}$ is equal to domain $d_\infty\lambda^{-1}_{d_\infty\Phi}$ of $\lambda^{-1}_{d_\infty\Phi}$, which is equal to domain $d_\infty\Phi$ of Φ. Similarly domain

$$d_\infty\rho_{c_\infty\Phi} \bullet_\infty (i_{s_\infty\Phi} *_\infty \Phi) \bullet_\infty \rho^{-1}_{d_\infty\Phi}$$

of composition $\rho_{c_\infty\Phi} \bullet_\infty (i_{s_\infty\Phi} *_\infty \Phi) \bullet_\infty \rho^{-1}_{d_\infty\Phi}$ is equal to domain $d_\infty\rho^{-1}_{d_\infty\Phi}$ of $\rho^{-1}_{d_\infty\Phi}$, which is equal to domain $d_\infty\Phi$ of Φ. We conclude, from this that relation 4 in the definition of equivalence relation R_∞ is compatible with domain function d_∞ of category F_∞. An analogous computation proves that

relation 4 in the definition of R_∞ is compatible with codomain function c_∞ of F_∞.

Let Φ, Ψ, Θ be a general morphisms in category F_∞. Suppose triple $\Phi, \Psi, \Theta$is compatible with respect to morphism functions of functors s_∞ and t_∞. In that case domain

$$d_\infty A_{c_\infty\Phi, c_\infty\Psi, c_\infty\Theta} \bullet_\infty [\Theta *_\infty (\Psi *_\infty \Phi)]$$

is equal to composition $A_{c_\infty\Phi, c_\infty\Psi, c_\infty\Theta} \bullet_\infty [\Theta *_\infty (\Psi *_\infty \Phi)]$ is equal to domain $d_\infty\Theta *_\infty (\Psi *_\infty \Phi)$ of composition $\Theta *_\infty (\Psi *_\infty \Phi)$ which in turn is equal to composition $d_\infty\Theta d_\infty\Psi d_\infty\Phi$ in $\mathcal{B}$. Now, domain

$$d_\infty[(\Theta *_\infty \Psi) *_\infty \Phi] \bullet_\infty A_{d_\infty\Phi, d_\infty\Psi, d_\infty\Theta}$$

is equal to domain $d_\infty A_{d_\infty\Phi, d_\infty\Psi, d_\infty\Theta}$ of associator $A_{d_\infty\Phi, d_\infty\Psi, d_\infty\Theta}$ associated to triple Φ, Ψ, Θ, which is, by definition, equal to composition $d_\infty\Theta d_\infty\Psi d_\infty\Phi$ in $\mathcal{B}$. We conclude that relation 5 in the definition of equivalence relation R_∞ is compatible with domain function d_∞ of category F_∞. An analogous computation proves that relation 5 in the definition of R_∞ is compatible with codomain function c_∞ of category F_∞.

Finally, the fact that equivalence relation R_∞ is compatible with respect to composition function $\bullet_\infty$ on category F_∞ follows from the fact that F_∞ is the limit in 2-category **Cat** of a sequence of free categories. ■

Definition 3.8. Let $\mathcal{B}$ be a decorated bicategory. We write V_∞ for quotient F_∞/R_∞ of category F_∞ associated to $\mathcal{B}$ modulo relation R_∞. We keep writing d_∞, c_∞, and $\bullet_\infty$ for domain, codomain, and composition operation functions in V_∞ respectively. We write H_∞ for collection of morphisms of category V_∞. Collection H_∞ is equal to the quotient E_∞/R_∞ of collection of morphisms E_∞ of category F_∞ modulo equivalence relation R_∞.

Lemma 3.9. *Let $\mathcal{B}$ be a decorated bicategory. In that case source and target functors s_∞ and t_∞, and horizontal composition function $*_\infty$ are all compatible with equivalence relation R_∞ defining category V_∞.*

Proof. Let $\mathcal{B}$ be a decorated bicategory. We wish to prove that in that case source and target functors s_∞ and t_∞, and horizontal composition function $*_\infty$ defined on category F_∞ associated to $\mathcal{B}$ are compatible with equivalence relation R_∞.

Let first Φ_i, Ψ_i, $i = 1, 2$ be morphisms in F_∞ such that pairs Φ_i, Ψ_i, $i = 1, 2$ are compatible with respect to domain and codomain functions d_∞

and c_∞ in F_∞ and such that pairs Φ_1, Φ_2 and Ψ_1, Ψ_1 are compatible with respect to source and target functors s_∞, t_∞ in F_∞. In that case source

$$s_\infty(\Psi_2 *_\infty \Psi_1) \bullet_\infty (\Phi_2 *_\infty \Phi_1)$$

of composition $(\Psi_2 *_\infty \Psi_1) \bullet_\infty (\Phi_2 *_\infty \Phi_1)$ is equal to composition

$$s_\infty(\Psi_2 *_\infty \Psi_1) s_\infty(\Phi_2 *_\infty \Phi_1)$$

in decoration $\mathcal{B}^*$ of $\mathcal{B}$ of vertical morphisms $s_\infty(\Phi_2 *_\infty \Phi_1)$ and $s_\infty(\Psi_2 *_\infty \Psi_1)$. Source $s_\infty(\Phi_2 *_\infty \Phi_1)$ of concatenation $\Phi_2 *_\infty \Phi_1$ is equal to source $s_\infty \Phi_1$ of Φ_1 and source $s_\infty(\Psi_2 *_\infty \Psi_1)$ of concatenation $\Psi_2 *_\infty \Psi_1$ is equal to source $s_\infty \Psi_2$. Source

$$s_\infty(\Psi_2 *_\infty \Psi_1) \bullet_\infty (\Phi_2 *_\infty \Phi_1)$$

of composition $(\Psi_2 *_\infty \Psi_1) \bullet_\infty (\Phi_2 *_\infty \Phi_1)$ is thus equal to composition $s_\infty \Psi_1 s_\infty \Phi_1$ in $\mathcal{B}^*$. Now source

$$s_\infty(\Psi_2 \bullet_\infty \Phi_2) *_\infty (\Psi_1 \bullet_\infty \Phi_1)$$

of concatenation $(\Psi_2 \bullet_\infty \Phi_2) *_\infty (\Psi_1 \bullet_\infty \Phi_1)$ is equal to source $s_\infty(\Psi_1 \bullet_\infty \Phi_1)$ of composition $\Psi_1 \bullet_\infty \Phi_1$. Now, source $s_\infty(\Psi_1 \bullet_\infty \Phi_1)$ is equal to composition $s_\infty \Psi_1 s_\infty \Phi_1$ of source $s_\infty \Phi_1$ and $s_\infty \Psi_1$ in $\mathcal{B}^*$. This proves that source functor s_∞ is compatible with respect to relation 1 in the definition of equivalence relation R_∞. An analogous argument proves that target functor t_∞ in F_∞ is compatible with respect to relation 1 in the definition of equivalence relation R_∞.

Let now Φ and Ψ be morphisms in $\mathbb{G}$ such that pair Φ, Ψ is compatible with respect to domain and codomain functions d_∞ and c_∞ in F_∞. Suppose first that Φ and Ψ are globular morphisms in $\mathcal{B}$ such that domain in $\mathcal{B}$ of domain and codomain of Φ and Ψ in $\mathcal{B}$ respectively are equal to object a in $\mathcal{B}$. In that case source $s_\infty \Psi \bullet_\infty \Phi$ of composition $\Psi \bullet_\infty \Phi$ is equal to composition $s_\infty \Psi s_\infty \Phi$ of $s_\infty \Phi$ and $s_\infty \Phi$ in $\mathcal{B}^*$. Now source $s_\infty \Psi$ of Φ and source $s_\infty \Psi$ of Ψ are both equal to the identity endomorphism id_a of object a in decoration $\mathcal{B}^*$ of $\mathcal{B}$. Now domain in $\mathcal{B}$ of vertical composition $\Psi \bullet \Phi$ in $\mathcal{B}$ is equal to domain of Φ in $\mathcal{B}$ and thus domain of domain in $\mathcal{B}$ of vertical composition $\Psi \bullet \Phi$ of Φ and Ψ is equal to object a of $\mathcal{B}$. It follows, from this, that source $s_\infty \Psi \bullet \Phi$ of globular morphism in $\mathcal{B}$ formed as the vertical composition $\Psi \bullet \Phi$ of Φ and Ψ in $\mathcal{B}$ is equal to identity endomorphism id_a of object a in $\mathcal{B}$. Source functor s_∞ in F_∞ is thus compatible with restriction to collection of globular morphisms of $\mathcal{B}$ of relation 2 in the definition of

equivalence relation R_∞. Suppose now that morphisms Φ and Ψ are formal horizontal identities i_α and i_β respectively, of composable pair of vertical morphisms α, β in $\mathcal{B}$. In this case source $s_\infty i_\beta \bullet_\infty i_\alpha$ of vertical composition $i_\beta \bullet_\infty i_\alpha$ of i_α and i_β is equal to composition $s_\infty i_\beta s_\infty i_\alpha$ of $s_\infty i_\alpha$ and i_β in $\mathcal{B}^*$. Source $s_\infty i_\alpha$ is equal to morphism α and source $s_\infty i_\beta$ of i_β is equal to morphism β. We conclude that source $s_\infty i_\beta \bullet_\infty i_\alpha$ of composition $i_\beta \bullet_\infty i_\alpha$ is equal to composition $\beta\alpha$ of α and β in $\mathcal{B}^*$. Finally, source $s_\infty i_{\beta\alpha}$ of formal horizontal identity $i_{\beta\alpha}$ of composition $\beta\alpha$ is equal to composition $\beta\alpha$. This proves that source functor s_∞ is compatible with restriction to collection of formal horizontal identities of relation 2 in the definition of equivalence relation R_∞. We conclude that source functor s_∞ is compatible with respect to relation 2 in teh definition of equivalence relation R_∞. An analogous argument proves that target functor t_∞ in F_∞ is compatible with respect to relation 2 in the definition of equivalence relation R_∞.

Let now Φ and Ψ be globular morphisms in $\mathcal{B}$ such that pair Φ, Ψ is compatible with respect to source and target functors s_∞, t_∞ in F_∞. Source $s_\infty \Psi *_\infty \Phi$ of concatenation $\Psi *_\infty \Phi$ of Φ and Ψ is equal to source $s_\infty \Phi$ which is equal to identity endomorphism in $\mathcal{B}^*$ of domain of domain in $\mathcal{B}$ of Φ. Now, source $s_\infty \Psi * \Phi$ of globular morphism in $\mathcal{B}$ formed as the horizontal composition $\Psi * \Phi$ in $\mathcal{B}$ of Φ and Ψ is equal to identity endomorphism in $\mathcal{B}^*$ of domain of domain in $\mathcal{B}$ of composition $\Psi * \Phi$. Domain of domain of horizontal composition $\Phi * \Phi$ is equal to domain of domain of Φ. We conclude that source $s_\infty \Psi * \Phi$ of horizontal composition $\Psi * \Phi$ is equal to identity endomorphism in $\mathcal{B}^*$ of domain of domain in $\mathcal{B}$ of Φ. Source functor s_∞ in $\mathcal{B}$ is thus compatible with relation 3 in the definition of equivalence relation R_∞. An identical argument proves that target functor t_∞ in F_∞ is compatible with relation 3 in the definition of equivalence relation R_∞.

Let now Φ be a general morphism in F_∞. In that case source

$$s_\infty \lambda_{c_\infty \Phi} \bullet_\infty (\Phi *_\infty i_{t_\infty \Phi}) \bullet_\infty \lambda^{-1}_{d_\infty \Phi}$$

of composition $\lambda_{c_\infty \Phi} \bullet_\infty (\Phi *_\infty i_{t_\infty \Phi}) \bullet_\infty \lambda^{-1}_{d_\infty \Phi}$ is equal to composition

$$s_\infty \lambda_{c_\infty \Phi} s_\infty (\Phi *_\infty i_{t_\infty \Phi}) s_\infty \lambda^{-1}_{d_\infty \Phi}$$

in $\mathcal{B}^*$ of $s_\infty \lambda_{c_\infty \Phi}$, $s_\infty (\Phi *_\infty i_{t_\infty \Phi})$, and $s_\infty \lambda^{-1}_{d_\infty \Phi}$. Now, since $\lambda_{c_\infty \Phi}$ and $\lambda_{d_\infty \Phi}$ are globular, composition

$$s_\infty \lambda_{c_\infty \Phi} s_\infty (\Phi *_\infty i_{t_\infty \Phi}) s_\infty \lambda^{-1}_{d_\infty \Phi}$$

is equal to source $s_\infty \Phi *_\infty i_{t_\infty \Phi}$ of $\Phi *_\infty i_{t_\infty \Phi}$, which is equal to source $s_\infty \Phi$ of Φ. We conclude that source functor s_∞ on F_∞ is compatible with relation 3 in the definition of equivalence relation R_∞. An analogous argument proves that target functor t_∞ is compatible with relation 4 in the definition of R_∞. Further, an analogous argument proves that source and target functors s_∞ and t_∞ in F_∞ are compatible with relation 5 in the definition of equivalence relation R_∞. ■

Observation 3.10. Let $\mathcal{B}$ be a decorated bicategory. By lemma 3.9 functors s_∞ and t_∞ descend to functors from category V_∞ to decoration $\mathcal{B}^*$ of $\mathcal{B}$. We keep denoting these functors by s_∞ and t_∞. Moreover, composition operation function $*_\infty$ descends to a function from collection $H_\infty \times_{\mathrm{Hom}_{\mathcal{B}^*}} H_\infty$ to collection H_∞ such that, by relation 1 in the definition of equivalence relation R_∞, together with composition operation function for horizontal morphisms in $\mathcal{B}$ forms a bifunctor from category $V_\infty \times_{\mathcal{B}^*} V_\infty$ to category V_∞. We denote this bifunctor by $*_\infty$.

Observation 3.11. Let $\mathcal{B}$ be a decorated bicategory. Let k be a positive integer. In that case relation R_∞ restricts to an equivalence relation in collection E_k associated to $\mathcal{B}$. We denote by H_k the quotient E_k/R_∞ of collection E_k modulo relation R_∞. Moreover, R_∞ restricts to an equivalence relation on collection of morphisms Hom_{F_k} of category F_k. Relation R_∞ is compatible with domain and codomain functions d_k and c_k of category F_k and is thus compatible with category structure of F_k. We denote by V_k the quotient F_k/R_∞ of category F_k modulo relation R_∞ and keep denoting by d_k, c_k, and $\bullet_k$ domain, codomain, and composition operation function in V_k. Functors s_{k+1} and t_{k+1} are compatible with R_∞ and thus induce functors from V_k to decoration $\mathcal{B}^*$ of $\mathcal{B}$. We keep denoting these functors by s_{k+1} and t_{k+1} respectively. Finally, observe that bifunction $*_k$ is compatible with relation R_∞ and thus defines a bifunction from collection of morphisms of category $V_k \times_{\mathcal{B}^*} V_k$ to collection of morphisms of V_k. This function, together with composition operation function for horizontal morphisms in $\mathcal{B}$ forms a bifunctor from $V_k \times_{\mathcal{B}^*} V_k$ to V_k. We keep denoting this bifunctor by $*_k$.

Observation 3.12. Let $\mathcal{B}$ be a decorated bicategory. Let m, k be positive integers such that m is less than or equal to k. In that case collection H_m is contained in collection H_k, category V_m is subcategory of category V_k, functors s_{m+1} and t_{m+1} are restrictions to category V_m of functors s_{k+1} and t_{k+1}, and bifunctor $*_m$ is restriction to $V_m \times_{\mathcal{B}^*} V_m$ of bifunctor $*_k$. Moreover, category V_∞ is equal to limit $\varinjlim V_k$ in 2-category **Cat** of sequence V_k, collection of morphisms H_∞ of V_∞ is equal to union $\bigcup_{k=1}^{\infty} H_k$ of sequence

H_k, and functors s_∞, t_∞ and $*_\infty$ are limits in 2-category **Cat**, of sequences of functors s_k, t_k and $*_k$ respectively.

Notation 3.13. Let $\mathcal{B}$ be a decorated bicategory. Pair formed by function associating horizontal identity id_a to every object a of $\mathcal{B}$ and function associating formal horizontal identity i_α to every vertical morphism α in $\mathcal{B}$ defines a functor from decoration $\mathcal{B}^*$ of $\mathcal{B}$ to category V_∞ associated to $\mathcal{B}$. We denote this functor by i_∞. For every positive integer k we denote codomain restriction to category V_k of functor i_∞ by i_k. Thus defined, i_k is a functor from decoration $\mathcal{B}^*$ of $\mathcal{B}$ to category V_k associated to $\mathcal{B}$ for every positive integer k.

Lemma 3.14. *Let $\mathcal{B}$ be a decorated bicategory. Let k be a positive integer. In that case functors s_{k+1}, t_{k+1}, i_k, and $*_k$ satisfy the following two conditions*

1. *The following two triangles commute*



2. *The following two squares commute, where q_1^k, q_2^k denote the left and right projection functors from $V_k \times_{\mathcal{B}^*} V_k$ to V_k respectively.*



Proof. Let $\mathcal{B}$ be a decorated bicategory. Let k be a positive integer. We wish to prove that conditions 1 and 2 above are satisfied.

We begin by proving that functors s_{k+1}, t_{k+1}, and i_k satisfy condition 1 above. Let α be a vertical morphism in $\mathcal{B}$. Formal horizontal identity i_α

associated to α, that is, the image $i_k\alpha$ of α under functor i_k, is a morphism in V_1. From this and from the fact that sequences s_{k+1} and t_{k+1} satisfy conditions of proposition 3.2 it follows that commutativity of triangles in 1 is equivalent to commutativity of triangles



which follows directly from the definition of functions d_1 and c_1. We now prove that functors s_{k+1}, t_{k+1}, and $*_k$ satisfy condition 2 above. Commutativity of squares in 2 when evaluated on morphisms in collection $H_k \times_{\mathrm{Hom}_{\mathcal{B}^*}} H_k$ of generators of $V_k \times_{\mathcal{B}^*} V_k$ follows from observation 3.4. General commutativity of squares in condition 2 follows from this and from the fact that all edges involved are functors. This concludes the proof of the lemma. ■

The following corollary follows directly from the previous lemma by taking limits.

Corollary 3.15. *Let $\mathcal{B}$ be a decorated bicategory. In that case functors s_∞, t_∞, i, and $*_\infty$ satisfy the following two conditions.*

1. *The following two triangles commute*



2. *The following two squares commute, where q_1^∞, q_2^∞ denote the left and right projection functors from $V_\infty \times_{\mathcal{B}^*} V_\infty$ to V_∞ respectively.*



Given a decorated bicategory $\mathcal{B}$ we will denote by $Q_{\mathcal{B}}$ pair formed by decoration $\mathcal{B}^*$ of $\mathcal{B}$ and category V_∞ associated to $\mathcal{B}$. The following theorem says that we can endow pair $Q_{\mathcal{B}}$ with the structure of a globularily generated double category.

Theorem 3.16. *Let $\mathcal{B}$ be a decorated bicategory. Pair $Q_{\mathcal{B}}$ together with functors s_∞, t_∞, i, bifunctor $*_\infty$, and collection of left and right identity transformations, and associator of $\mathcal{B}$, is a double category. Moreover, with this structure, double category $Q_{\mathcal{B}}$, is globularily generated.*

Proof. Let $\mathcal{B}$ be a decorated bicategory. We wish to prove in this case that pair $Q_{\mathcal{B}}$, together with functors s_∞, t_∞, i, bifunctor $*_\infty$, and collection of left and right identity transformations and associator of $\mathcal{B}$ is a globularily generated double category.

Formal horizontal identity functor i and horizontal composition bifunctor $*_\infty$ are compatible with functors s_∞ and t_∞ by corollary 3.15 Collections of left and right identity transformations of $\mathcal{B}$ form a natural transformation from functor $*_\infty(is_\infty \times id_{V_\infty})$ to identity endofunctor id_{V_∞} of V_∞ and a natural transformation from $*_\infty(id_{V_\infty} \times it_\infty)$ to identity endobifunctor id_{V_∞} of V_∞ respectively by the fact that morphisms in V_∞ satisfy relation 4 in the definition of relation R_∞ associated to $\mathcal{B}$. Collection of associators of $\mathcal{B}$ forms a natural transformation from composition $*_\infty(*_\infty \times id_{V_\infty})$ to composition $*_\infty(id_{V_\infty} \times *_\infty)$ by the fact that morphisms in V_∞ satisfy relation 5 in the definition of R_∞. Left and right identity and associator relations for pair $Q_{\mathcal{B}}$ again follow from the fact that morphisms in V_∞ satisfy relations 4 and 5 in the definition of R_∞. The fact that pair formed by functor i and bifunctor $*$ satisfies Mc Lane's triangle and pentagon relations with respect to left and right identity transformations and associator follows from the fact that components of left and right identity transformations and associator satisfy McLane's axioms for bicategory $\mathcal{B}$. This proves that $Q_{\mathcal{B}}$ with the structure described is a double category. A straightforward induction argument proves

that for every positive integer k every morphism of category V_k is globularily generated in $Q_{\mathcal{B}}$, from which it follows that double category $Q_{\mathcal{B}}$ is globularily generated. This concludes the proof of the theorem. ■

Definition 3.17. Let $\mathcal{B}$ be a decorated bicategory. We call globularily generated double category $Q_{\mathcal{B}}$ the free globularily generated double category associated to $\mathcal{B}$.

When free globularily generated double categories of different decorated bicategories are being considered we will write each decorated bicategory as a superindex for all the structure involved in free globularily generated double category associated to it e.g. We will write $s_\infty^{\mathcal{B}}, t_\infty^{\mathcal{B}}, *_\infty^{\mathcal{B}}, \bullet_\infty^{\mathcal{B}}, i^{\mathcal{B}}$ for double category structure of free double category $Q_{\mathcal{B}}$ associated to a decorated bicategory $\mathcal{B}$. The same convention will apply to every piece of structure involved in the free globularily generated double category construction. Notational conventions introduced in results and proofs of results in this section will be used throughout the paper without further reference.

Examples

Finally, we present globularily generated double category construction performed in a couple of examples of decorated bicategories.

Trivially decorated bicategories: Let B be a bicategory. We consider B as a trivially decorated bicategory. In that case the only vertical morphisms in B are identity endomorphisms. It follows that every formal horizontal identity in free globularily generated double category Q_B associated to B is a globular morphism in B. From this and from the fact that collection of globular morphisms in B is clearly closed under vertical and horizontal composition operations in Q_B it follows that free globularily generated double category Q_B associated to B is equal to B. Observe that in this case for every positive integer k collection H_k^B is equal to collection H_1^B and category V_k^B is equal to category V_1^B. Finally, observe that in this case decorated horizontalization H^*Q_B of free globularily generated double category Q_B associated to B is equal to B, that is, free globularily generated double category Q_B associated to trivially decorated bicategory B is a verticalization of B.

Decorated bicategory $2m\mathbb{Z}/2\mathbb{Z}_{\mathbb{Z}/2\mathbb{Z}}$**:** We write $m\mathbb{Z}/2\mathbb{Z}$ for single object monoidal category associated to group $\mathbb{Z}/2\mathbb{Z}$. Category $m\mathbb{Z}/2\mathbb{Z}$ has a single

object whose monoid of endomorphisms is $\mathbb{Z}/2\mathbb{Z}$. Composition and tensor product of morphisms in $m\mathbb{Z}/2\mathbb{Z}$ are both given by product operation in $\mathbb{Z}/2\mathbb{Z}$. Exchange property follows, in this case from commutativity of product operation in $\mathbb{Z}/2\mathbb{Z}$. We write $\mathcal{B}$ for decorated bicategory $2m\mathbb{Z}/2\mathbb{Z}_{\mathbb{Z}/2\mathbb{Z}}$ associated to monoidal category $m\mathbb{Z}/2\mathbb{Z}$, decorated by $m\mathbb{Z}/2\mathbb{Z}$. Decorated bicategory $\mathcal{B}$ has in this case a single object a, two vertical morphisms which we denote by 1 and -1, 1 denoting the identity endomorphism of a, and a single horizontal morphism, the horizontal identity endomorphism id_a of a. Decorated bicategory $\mathcal{B}$ has two globular morphisms which we denote by **1** and -**1**, **1** denoting the identity endomorphism of horizontal identity endomorphism id_a of a. Collection $E_1^{\mathcal{B}}$ associated to $\mathcal{B}$ thus has three morphisms, namely **1**, -**1**, and formal horizontal identity i_{-1} of vertical morphism -1. Category $V_1^{\mathcal{B}}$ has horizontal identity id_a of object a as single object with free product $\mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/2\mathbb{Z}$ as monoid of endomorphisms, with globular morphism -**1** and formal horizontal identity id_{-1} of -1 as generators. Now, endomorphism $(-\mathbf{1}) \bullet_\infty i_{-1} \bullet_\infty (-\mathbf{1})$ of horizontal identity endomorphism id_{-1} of -1 is globular in free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ but it is not a globular morphism in $\mathcal{B}$. This proves, on the one hand that collections $H_1^{\mathcal{B}}$ and $H_1^{Q_{\mathcal{B}}}$ are different and thus horizontal filtration of $Q_{\mathcal{B}}$ and free horizontal filtration associated to $\mathcal{B}$ are different, and on the other hand that free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ is not a verticalization of $\mathcal{B}$.

4 Canonical double projection

In this section we present the canonical double projection construction. We associate a double functor from free globularily generated double category associated to a decorated bicategory to any of its globularily generated verticalizations. We prove that this double functor is uniquely determined by a certain set of properties and that its morphism functor is complete. We interpret this by saying that canonical double projection canonically provides every globularily generated verticalization of a decorated bicategory with the structure of quotient of its free globularily generated double category. The following is the main theorem of this section.

Theorem 4.1. *Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. In that case there exists a unique double functor $\pi^C : Q_{\mathcal{B}} \to C$ such that restriction of decorated horizontalization $H^*\pi^C$ of π^C to $\mathcal{B}$ is equal to identity decorated*

endobifunctor $id_{\mathcal{B}}$ *of* $\mathcal{B}$*. Morphism functor of double bifunctor* π^C *is complete.*

Given a globularily generated verticalization C of a decorated bicategory $\mathcal{B}$ we call double functor π^C defined in the previous proposition, the canonical double projection associated to C. We divide the proof of theorem 4.1 in several steps. We begin by proving the existence part of theorem 3.16. We proceed in a similar way as we did with the free globularily generated double category construction in section 3.

Existence

Notation 4.2. Let C be a double category. We denote by q^C function from collection of evaluations $\mathrm{Hom}_{C_1{}_{s,t}}$ of collection of 2-morphisms Hom_{C_1} of C with respect to morphism functions of source and target functors s, t of C to collection of 2-morphisms Hom_{C_1} of C such that for every evaluation Φ of a compatible sequence $\Psi_1, ..., \Psi_k$ of 2-morphisms in C, the image $q^C\Phi$ of Φ under q^C equals horizontal composition in C of 2-morphisms $\Psi_1, ..., \Psi_k$ following parenthesis pattern defining Φ.

Lemma 4.3. *Let* $\mathcal{B}$ *be a decorated bicategory. Let* C *be a globularily generated double category. Suppose* C *is a verticalization of* $\mathcal{B}$*. In that case there exists a pair, formed by a sequence of functions* $E_k^{\pi^C} : E_k^{\mathcal{B}} \to Hom_{C_1}$ *and a sequence of functors* $F_k^{\pi^C} : F_k^{\mathcal{B}} \to C_1$*, with* k *running through collection of all positive integers, such that the following conditions are satisfied*

1. *Restriction of function* $E_1^{\pi^C}$*, to collection* $\mathbb{G}^{\mathcal{B}}$ *of globular morphisms of* $\mathcal{B}$ *and horizontal identities of vertical morphisms of* $\mathcal{B}$*, is equal to the identity function* $id_{\mathbb{G}^{\mathcal{B}}}$ *of* $\mathbb{G}^{\mathcal{B}}$*.*

2. *For every pair of positive integers* m, k *such that* m *is less than or equal to* k *restriction of function* $E_k^{\pi^C}$ *to collection of morphisms of category* $F_m^{\mathcal{B}}$ *is equal to morphism function of functor* $F_m^{\pi^C}$*, and restriction to collection* $E_m^{\mathcal{B}}$ *of morphism function of functor* $F_k^{\pi^C}$ *is equal to function* $E_m^{\pi^C}$*.*

3. *The following two triangles commute for every positive integer* k



4. The following two triangles commute for every positive integer k



5. The following two triangles commute for every positive integer k



6. The following square commutes for every positive integer k



Moreover, conditions 1-5 above determine pair of sequences $E_k^{\pi^C}$ *and* $F_k^{\pi^C}$.

Proof. Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. We wish to construct a sequence of functions $E_k^{\pi^C}$ from collection $E_k^{\mathcal{B}}$ to collection of morphisms Hom_{C_1} of morphism category C_1 of C, and a sequence of functors $F_k^{\pi^C}$ from category $F_k^{\mathcal{B}}$ to morphism category C_1 of C with k running through collection of positive integers, in such a way that pair of sequences $E_k^{\pi^C}$ and $F_k^{\pi^C}$ satisfies conditions 1-6 of the lemma.

We proceed inductively on k. We begin with the definition of function $E_1^{\pi^C}$. Observe first that from the fact that C is a verticalization of decorated bicategory $\mathcal{B}$ it follows that collection of vertical morphisms of $\mathcal{B}$ is equal to collection of vertical morphisms of C. There is thus an obvious identification between collection of formal horizontal identities of vertical morphisms of $\mathcal{B}$ and collection of horizontal identities of vertical morphisms of C. We use this identification and consider collection of formal horizontal identities of vertical morphisms of $\mathcal{B}$ and collection of horizontal identities of vertical morphisms of C as being the same. Observe that the fact that C is a verticalization of $\mathcal{B}$ also implies that collection of globular morphisms of $\mathcal{B}$ is equal to collection of globular morphisms of C. Collection $\mathbb{G}^{\mathcal{B}}$ defining collection $E_1^{\mathcal{B}}$ of morphisms of category $V_\infty^{\mathcal{B}}$ is thus equal to collection of fundamental generators $\mathbb{G}^C$ of globularily genrated double category C. We make function $E_1^{\pi^C}$ to be equal to composition $q^C \mu_{id_{\mathbb{G}^{\mathcal{B}}}, id_{\mathcal{B}^*}}$ of function $\mu_{id_{\mathbb{G}^{\mathcal{B}}}, id_{\mathcal{B}^*}}$ associated to pair $id_{\mathbb{G}^{\mathcal{B}}}, id_{\mathcal{B}^*}$ and function q^C associated to double category C. Thus defined $E_1^{\pi^C}$ is a function from collection $E_1^{\mathcal{B}}$ to collection of morphisms Hom_{C_1} of morphism category C_1 of C. Moreover, from the way it was defined, and from ? it follows that function $E_1^{\pi^C}$ satisfies condition 1 and conditions 3-5 of the lemma. We now define functor $F_1^{\pi^C}$ as follows: Observe first that from the fact that C is a verticalization of $\mathcal{B}$ it follows that collection of horizontal morphisms of $\mathcal{B}$ is equal to collection of horizontal morphisms of C. We make object function of functor $F_1^{\pi^C}$ to be equal to identity function $id_{\mathcal{B}_1}$ of collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$. From the fact that function $E_1^{\pi^C}$ satisfies condition 3 of the lemma and from the fact that $E_1^{\mathcal{B}}$ freely generates $F_1^{\mathcal{B}}$ with respect to pair of functions $d_1^{\mathcal{B}}, c_1^{\mathcal{B}}$ it follows that there exists a unique extension of $E_1^{\pi^C}$ to a functor from category $F_1^{\mathcal{B}}$ to morphism category C_1 of C. We make $F_1^{\pi^C}$ to be this extension. Thus defined $F_1^{\pi^C}$ trivially satisfies condition 2 of the lemma with respect to function $E_1^{\pi^C}$. The fact that functor $F_1^{\pi^C}$ satisfies condition 5 of the lemma follows from the fact that function $E_1^{\pi^C}$ satisfies condition 4 and from functoriality of s_2

and s.

Let k be a positive integer strictly greater than 1. Assume now that for every positive integer m strictly less than k function $E_m^{\pi^C}$ from collection $E_m^{\mathcal{B}}$ to collection of morphisms Hom_{C_1} of morphism category C_1 of C and functor $F_m^{\pi^C}$ from category $F_m^{\mathcal{B}}$ to morphism category C_1 of C have been defined, in such a way that pair of sequences $E_m^{\pi^C}$ and $F_m^{\pi^C}$ with m running through collection of positive integers strictly less than k satisfies conditions 1-6 of the lemma. We now construct function $E_k^{\pi^C}$ from collection $E_k^{\mathcal{B}}$ to collection of morphisms Hom_{C_1} of category of morphisms C_1 of C and functor $F_k^{\pi^C}$ from category $F_k^{\mathcal{B}}$ to category of morphisms C_1 of C such that pair $E_k^{\pi^C}$ and $F_k^{\pi^C}$ satisfies conditions 1-6 of the lemma with respect to pair of sequences $E_m^{\pi^C}$ and $F_m^{\pi^C}$ with m running through collection of positive integers strictly less than k.

We first define function $E_k^{\pi^C}$. Denote first From the assumption that functor $F_{k-1}^{\pi^C}$ satisfies condition 5 of the lemma it follows that function $\mu_{F_{k-1}^{\pi^C},id_{\mathcal{B}^*}}$ is well defined. We make $E_k^{\pi^C}$ to be composition $q^C\mu_{F_{k-1}^{\pi^C},id_{\mathcal{B}^*}}$ of function $\mu_{F_{k-1}^{\pi^C},id_{\mathcal{B}^*}}$ associated to pair $F_{k-1}^{\pi^C}, id_{\mathcal{B}^*}$ and function q^C associated to C. Thus defined $E_k^{\pi^C}$ is a function from collection $E_k^{\mathcal{B}}$ to collection of morphisms Hom_{C_1} of morphism category C_1 of C. From the way it was defined it is clear that function $E_k^{\pi^C}$ satisfies conditions 4 and 6 of the lemma. From induction hypothesis it follows that $E_k^{\pi^C}$ satisfies conditions 1 and 2. Function $E_k^{\pi^C}$ satisfies condition 3 of the lemma by the fact that it satisfies condition 2 and by functoriality of $F_{k-1}^{\pi^C}$. We now define functor $F_k^{\pi^C}$. By the fact that function $E_k^{\pi^C}$ satisfies condition 3 of the lemma it follows that there is a unique extension of $E_k^{\pi^C}$ to a functor from category $F_k^{\mathcal{B}}$ to category C_1. We make $F_k^{\pi^C}$ to be this functor. Thus defined $F_k^{\pi^C}$ satisfies condition 2 of the lemma. This follows from the way $F_k^{\pi^C}$ was constructed and from the fact that condition 2 is already satisfied by function $E_k^{\pi^C}$. From the fact that $E_k^{\pi^C}$ satisfies condition 4 it follows that functor $F_k^{\pi^C}$ satisfies condition 5 of the lemma. We have thus constructed, recursively, a pair of sequences $E_k^{\pi^C}$ and $F_k^{\pi^C}$ satisfying the conditions of the lemma. This concludes the proof. ■

Observation 4.4. Let $\mathcal{B}$ be a decorated bicategory. Condition 2 of lemma 4.3 implies that for every pair of positive integers m, k such that m is less than or equal to k restriction of function $E_k^{\pi^C}$ to collection $E_m^{\mathcal{B}}$ is equal to function $E_m^{\pi^C}$ and that restriction of functor $F_k^{\pi^C}$ to category $F_m^{\mathcal{B}}$ is equal to

functor $F_m^{\pi^C}$.

Notation 4.5. Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. In that case we write $E_\infty^{\pi^C}$ for the limit $\varinjlim E_k^{\pi^C}$ in **Set** of sequence $E_k^{\pi^C}$. Thus defined $E_\infty^{\pi^C}$ is a function from collection $E_\infty^{\mathcal{B}}$ associated to $\mathcal{B}$ to collection of 2-morphisms Hom_{C_1} of C. Moreover, we write $F_\infty^{\pi^C}$ for the limit $\varinjlim F_k^{\pi^C}$ in 2-category **Cat** of sequence $F_k^{\pi^C}$. Thus defined $F_\infty^{\pi^C}$ is a functor from category $F_\infty^{\mathcal{B}}$ to morphism category C_1 of double category C. Function $E_\infty^{\pi^C}$ is equal to morphism function of functor $F_\infty^{\pi^C}$.

The following lemma follows directly from lemma 4.3 and observation 4.4.

Lemma 4.6. *Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. In that case functions $E_\infty^{\pi^C}$ and $F_\infty^{\pi^C}$ associated to C satisfy the following conditions*

1. *For every positive integer k restriction of function $E_\infty^{\pi^C}$ to collection of morphisms of category $F_k^{\mathcal{B}}$ is equal to function $E_k^{\pi^C}$. Restriction of functor $F_k^{\pi^C}$ to category $F_k^{\mathcal{B}}$ is equal to functor $F_k^{\pi^C}$.*

2. *The following two triangles commute*



3. *The following triangle commutes*



Lemma 4.7. *Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. In that case functor $F_\infty^{\pi^C}$ associated to C is compatible with equivalence relation R_∞.*

Proof. Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. We wish to prove that functor $F_\infty^{\pi^C}$ associated to C is compatible with equivalence relation R_∞ defined on collection of morphisms $E_\infty^{\mathcal{B}}$ of $F_\infty^{\mathcal{B}}$. Compatibility of functor $F_\infty^{\pi^C}$ with respect to relation 1 in the definition of equivalence relation R_∞ follows from functoriality of $F_\infty^{\pi^C}$ together with the fact that $F_\infty^{\pi^C}$ satisfies conditions 5 and 6 of lemma 3.2.

We now prove that functor $F_\infty^{\pi^C}$ is compatible with relation 2 in the definition of R_∞. Let first Φ and Ψ be globular morphisms of $\mathcal{B}$ such that pair Φ, Ψ is compatible with respect to domain and codomain functions d_∞ and c_∞ of $F_\infty^{\mathcal{B}}$. In that case the image $F_\infty^{\pi^C}\Psi \bullet_\infty \Phi$ of vertical composition $\Psi \bullet_\infty \Phi$ of Φ and Ψ under functor $F_\infty^{\pi^C}$ is equal to the image $F_1^{\pi^C}\Psi \bullet_\infty \Psi$ of composition $\Psi \bullet_\infty \Phi$ under functor $F_1^{\pi^C}$, which is, by functoriality of $F_1^{\pi^C}$ equal to composition $F_1^{\pi^C}\Psi \bullet F_1^{\pi^C}\Phi$ in C of $F_1^{\pi^C}\Phi$ and $F_1^{\pi^C}\Psi$, which is equal, by the definition of $F_1^{\pi^C}$ to vertical composition $\Psi \bullet \Phi$ of Φ and Ψ in C. Now, the image $F_\infty^{\pi^C}\Psi \bullet \Phi$ of vertical composition $\Psi \bullet \Phi$ in $\mathcal{B}$ of Φ and Ψ under functor $F_\infty^{\pi^C}$ is equal to the image $E_1^{\pi^C}\Psi \bullet \Phi$ of composition $\Psi \bullet \Phi$ under function $E_1^{\pi^C}$, which is equal, by the way function $E_1^{\pi^C}$ was defined, to vertical composition $\Psi \bullet \Phi$ of Φ and Ψ in C. Functor $F_\infty^{\pi^C}$ is thus compatible with respect to relation 2 in the definition of R_∞ when restricted to globular morphisms of $\mathcal{B}$. Let now α and β be vertical morphisms of $\mathcal{B}$ such that pair α, β is composable in decoration $\mathcal{B}^*$ of $\mathcal{B}$. In that case the image $F_\infty^{\pi^C} i_\beta \bullet_\infty i_\alpha$ of vertical composition $i_\beta \bullet_\infty i_\alpha$ of horizontal identities i_α and i_β of α and β under functor $F_\infty^{\pi^C}$ is equal to image $F_1^{\pi^C} i_\beta \bullet_\infty i_\alpha$ of composition $i_\beta \bullet_\infty i_\alpha$ under functor $F_1^{\pi^C}$, which is equal to vertical composition $F_1^{\pi^C} i_\beta \bullet F_1^{\pi^C} i_\beta$, in C, of $F_1^{\pi^C} i_\alpha$ and $F_1^{\pi^C} i_\beta$. This is equal, again by the definition of $F_1^{\pi^C}$, to vertical composition $i_\beta \bullet i_\alpha$ of i_α and i_β. Now, image $F_\infty^{\pi^C} i_{\beta\alpha}$ of formal horizontal identity $i_{\beta\alpha}$ of composition $\beta\alpha$ of α and β under functor $F_\infty^{\pi^C}$ is equal to image $E_1^{\pi^C} i_{\beta\alpha}$ of $i_{\beta\alpha}$ under function $E_1^{\pi^C}$ which is, by the way function $E_1^{\pi^C}$ was defined, equal to horizontal identity $i_{\beta\alpha}$ of $\beta\alpha$ in C, that is, image $F_\infty^{\pi^C} i_{\beta\alpha}$ of $i_{\beta\alpha}$ under functor $F_\infty^{\pi^C}$ is equal to vertical composition $i_\beta \bullet i_\alpha$ of i_α and i_β in C. We conclude that functor $F_\infty^{\pi^C}$ is compatible with relation 2 in the definition of equivalence relation R_∞ when restricted to formal horizontal identities and thus functor $F_\infty^{\pi^C}$ is compatible with relation

2 in the definition of R_∞.

We now prove that functor $F_\infty^{\pi^C}$ is compatible with relation 3 in the definition of R_∞. Let Φ and Ψ be globular morphisms in $\mathcal{B}$ such that pair Φ, Ψ is compatible with respect to functors s_∞ and t_∞. In that case image $F_\infty^{\pi^C} \Psi *_\infty \Phi$ of concatenation $\Psi *_\infty \Phi$ under functor $F_\infty^{\pi^C}$ is equal to image $E_1^{\pi^C} \Psi *_1 \Phi$ of concatenation $\Psi *_1 \Phi$ in $E_1^{\mathcal{B}}$ of Φ and Ψ under function $E_1^{\pi^C}$. This is equal, by the fact that function $E_1^{\pi^C}$ satisfies condition 6 of lemma 3.2, to horizontal composition $E_1^{\pi^C} \Psi * E_1^{\pi^C} \Phi$ of $E_1^{\pi^C} \Phi$ and $E_1^{\pi^C} \Psi$ in C, which, by the definition of $E_1^{\pi^C}$ is equal to horizontal composition $\Psi * \Phi$ of Φ and Ψ in $\mathcal{B}$. Now, image $F_\infty^{\pi^C} \Psi * \Psi$ of horizontal composition $\Psi * \Phi$ of Φ and Ψ in $\mathcal{B}$ under functor $F_\infty^{\pi^C}$ is equal to image $E_1^{\pi^C} \Phi * \Psi$ of horizontal composition $\Psi * \Phi$ in $\mathcal{B}$ under function $E_1^{\pi^C}$. This is equal, again by the way function $E_1^{\pi^C}$ was defined, to horizontal composition $\Psi * \Phi$ of Φ and Ψ in $\mathcal{B}$. We conclude that functor $F_\infty^{\pi^C}$ is compatible with relation 3 in the definition of equivalence relation R_∞.

Finally, compatibility of functor $F_\infty^{\pi^C}$ with relations 4 and 5 in the definition of relation R_∞ follows from conditions 3 and 5 of proposition 3.2 and from the fact that the identity decorated endobifunctor $id_{\mathcal{B}}$ of $\mathcal{B}$ carries left and right identity transformations to left and right identity transformations and associators to associators. This concludes the proof of the lemma.

■

Notation 4.8. Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. We will write $V_\infty^{\pi^C}$ for functor from category $V_\infty^{\mathcal{B}}$ to morphism category C_1 of C induced by functor $F_\infty^{\pi^C}$ and equivalence relation $R_\infty^{\mathcal{B}}$. Further, we write $H_\infty^{\pi^C}$ for morphism function of functor $V_\infty^{\pi^C}$.

The proof of the following lemma follows directly from lemma 4.6 by taking limits.

Lemma 4.9. *Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. In that case functor $V_\infty^{\pi^C}$ satisfies the following conditions.*

1. *The following two triangles commute*



2. The following triangle commutes for every positive integer k



We now prove the existence part of theorem 1.

Proof: Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. We wish to construct a double functor $\pi^C : Q_{\mathcal{B}} \to C$ from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to C such that decorated horizontalization $H^*\pi^C$ is equal to identity decorated endobifunctor $id_{\mathcal{B}}$ of $\mathcal{B}$.

We make double functor π^C associated to C to be equal to pair $(id_{\mathcal{B}^*}, V_\infty^{\pi^C})$ formed by identity endofunctor $id_{\mathcal{B}^*}$ of decoration $\mathcal{B}^*$ of $\mathcal{B}$ and functor $V_\infty^{\pi^C}$ constructed above. Pair π^C is a double functor from $Q_{\mathcal{B}}$ to C by lemma 4.9 and by the fact that it clearly intertwinesformal identity functor i_∞ in $Q_{\mathcal{B}}$ and identity functor i in C. The fact that restriction of decorated horizontalization $H^*\pi^C$ of double functor π^C to collection $\mathbb{G}^{\mathcal{B}}$ is equal to identity decorated endobifunctor $id_{\mathcal{B}}$ of $\mathcal{B}$ follows directly from the way functor $V_\infty^{\pi^C}$ was defined. This concludes the proof. ■

Definition 4.10. Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. We call double functor π^C defined in the above proof the canonical double projection associated to globularily generated verticalization C of $\mathcal{B}$.

Fullness

We now prove fullness part of theorem 1. We begin with the following lemma.

Lemma 4.11. *Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. Let k be a positive integer. In that case the image, under morphism function $H_\infty^{\pi^C}$ of functor $V_\infty^{\pi^C}$, of collection $H_k^{\mathcal{B}}$ is equal to k-th term H_k^C of horizontal filtration of C. Moreover, the image category of restriction to category $V_k^{\mathcal{B}}$ of functor $V_\infty^{\pi^C}$ is equal to k-th term V_k^C of vertical filtration associated to C.*

Proof. Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. Let k be a positive integer. We wish to prove that the image, under morphism function $H_\infty^{\pi^C}$ of canonical projection π^C associated to C, of subcollection $H_k^{\mathcal{B}}$ of collection of morphisms $H_\infty^{\mathcal{B}}$ of $Q_{\mathcal{B}}$ is equal to k-th term H_k^C of horizontal filtration of C. Moreover, we wish to prove that the image category of restriction to subcategory $V_k^{\mathcal{B}}$ of morphism category $V_\infty^{\mathcal{B}}$, of free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$, of morphism functor $V_\infty^{\pi^C}$ of canonical projection π^C associated to C, is equal to k-th term V_k^C of vertical filtration associated to C. We proceed by induction on k.

We prove first that the image, under function $H_\infty^{\pi^C}$, of collection $H_1^{\mathcal{B}}$ is equal to the first term H_1^C of horizontal filtration of C. From the fact that collection $H_1^{\mathcal{B}}$ is contained in first term $H_1^{Q_{\mathcal{B}}}$ of horizontal filtration of free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$, and from the fact that π^C is a double functor, it follows that the image, under $H_\infty^{\mathcal{B}}$, of collection $H_1^{\mathcal{B}}$ is contained in first term H_1^C of horizontal filtration associated to C. Now, function $H_\infty^{\pi^C}$ acts as the identity function when restricted to collection $\mathbb{G}^{\mathcal{B}}$ of globular morphisms of $\mathcal{B}$ and horizontal identities of vertical morphisms of $\mathcal{B}$. It follows, from this, from the fact that function $H_\infty^{\pi^C}$ satisfies condition 2 of lemma 4.9, and from the fact that collection H_1^C is collection of horizontal composition of elements of $\mathbb{G}^{\mathcal{B}}$, that the image of collection $H_1^{\mathcal{B}}$ under $H_\infty^{\mathcal{B}}$ contains collection H_1^C. We conclude that the image, under morphism function $H_\infty^{\pi^C}$ of morphism functor $V_\infty^{\pi^C}$ of canonical projection π^C associated to C, of collection $H_1^{\mathcal{B}}$ of morphisms of $Q_{\mathcal{B}}$ is equal to first term H_1^C of horizontal filtration of C. We now prove that the image category of category $V_1^{\mathcal{B}}$ associated to $\mathcal{B}$, under morphism functor $V_\infty^{\pi^C}$ of projection π^C, is equal to first term V_1^C of vertical filtration of C. From the previous argument, and the fact that functor $V_\infty^{\pi^C}$ satisfies condition 2 of lemma 4.9 it follows that the image, under functor $V_\infty^{\pi^C}$, of collection $H_1^{\mathcal{B}}$

of generators of category $V_1^{\mathcal{B}}$, is equal to collection H_1^C of generators of first term V_1^C of vertical filtration of C. This, together with the fact that $V_\infty^{\pi^C}$ is a functor, implies that the image category of $V_1^{\pi^C}$ under functor $V_\infty^{\pi^C}$ is precisely V_1^C.

Let now k be a positive integer strictly greater than 1. Suppose that for every positive integer m strictly less than k, the image of collection $H_m^{\mathcal{B}}$ under function $H_\infty^{\pi^C}$ is equal to collection H_m^C and that image category of category $V_m^{\mathcal{B}}$, under functor $V_\infty^{\pi^C}$, is equal to category V_m^C. We now prove that the image, under function $H_\infty^{\pi^C}$, of collection $H_k^{\mathcal{B}}$ is precisely k-th term H_k^C of horizontal filtration of C. From the fact that collection $H_k^{\mathcal{B}}$ is contained in the k-th term $H_k^{Q_{\mathcal{B}}}$ of horizontal filtration of $Q_{\mathcal{B}}$ and from the fact that π^C is a double functor it follows that the image, under function $H_\infty^{\pi^C}$ is contained in k-th term H_k^C of horizontal filtration of C. Now, since function $H_\infty^{\pi^C}$ satisfies condition 3 of proposition 4.6, induction hypothesis implies that image, under $H_\infty^{\pi^C}$, of collection of morphisms $\mathrm{Hom}_{V_{k-1}^{\mathcal{B}}}$ of category $V_{k-1}^{\mathcal{B}}$ is precisely collection of morphisms $\mathrm{Hom}_{V_{k-1}^C}$ of $k-1$-th term V_{k-1}^C of vertical filtration of C. It follows, from this, from the fact that $H_\infty^{\pi^C}$ satisfies condition 3 of lemma 4.3 and from the fact that every morphism in H_k^C is the horizontal composition of a composable sequence of morphisms in $\mathrm{Hom}_{V_{k-1}^C}$ that the image, under function $H_\infty^{\pi^C}$, of collection $H_k^{\mathcal{B}}$ contains collection H_k^C. We thus conclude that image, under function $H_\infty^{\pi^C}$, of collection $H_k^{\mathcal{B}}$ is equal to k-th term H_k^C of horizontal filtration of C. Finally, we prove that image category, under morphism functor $V_\infty^{\pi^C}$ of π^C, of category $V_k^{\mathcal{B}}$ is precisely k-th term V_k^C of vertical filtration associated to C. From the the previous argument and from the fact that functor $V_\infty^{\pi^C}$ satisfies condition ? of proposition ? it follows that image of collection $H_k^{\mathcal{B}}$ of generators of category $V_k^{\mathcal{B}}$ under functor $V_\infty^{\pi^C}$ is equal to collection H_k^C of generators of category V_k^C. This, together with functoriality of $V_k^{\pi^C}$ implies that image category of restriction, to category $V_k^{\mathcal{B}}$, of morphism functor $V_\infty^{\pi^C}$ of canonical projection π^C associated to C, is equal to n-th term V_k^C of vertical filtration of C. This concludes the proof. ■

We now prove the fullness part of theorem 1.

Proof: Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Suppose C is a verticalization of $\mathcal{B}$. We wish to prove, in this case, that morphism functor $V_\infty^{\pi^C}$ of canonical projection π^C associated

to C is complete.

Let k be a positive integer. In that case restriction, to subcategory $V_k^{\mathcal{B}}$ of morphism category $V_\infty^{\mathcal{B}}$ of free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$, of functor $V_\infty^{\pi^C}$ defines, by lemma 4.11, a functor from category $V_k^{\mathcal{B}}$ to k-th term V_k^C of vertical filtration of double category C. We denote this functor by $\tilde{V}_k^{\pi^C}$. The fact that functor $V_\infty^{\pi^C}$ satisfies condition 1 of lemma 4.6 implies that for every pair of integers m, k such that m is less than or equal to k, functor $\tilde{V}_m^{\pi^C}$ is equal to restriction, to category $\tilde{V}_m^{\mathcal{B}}$, of functor $\tilde{V}_k^{\pi^C}$. Sequence of functors $\tilde{V}_k^{\pi^C}$ is thus a direct system in 2-category **Cat**. Functor $V_\infty^{\pi^C}$ is equal to its limit $\varinjlim \tilde{V}_k^{\pi^C}$ in **Cat** of direct system $\tilde{V}_k^{\pi^C}$. This, together with the fact, following lemma 4.11, that functor $\tilde{V}_k^{\pi^C}$ is complete for every positive integer k completes the proof of the proposition. ■

We interpret this part of theorem 1 by saying that although free globularily generated double category $Q_{\mathcal{B}}$ of a given decorated bicategory $\mathcal{B}$ may not always be a verticalization of $\mathcal{B}$, it projects onto every globularily generated verticalization of $\mathcal{B}$. We now prove the uniqueness part of theorem 1.

Uniqueness

We begin the proof of uniqueness part of theorem 1 by extending notation used in the proof of completeness part of theorem 1 above.

Notation 4.12. Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Let $T : Q_{\mathcal{B}} \to C$ be a double functor from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to C. Let k be a positive integer. In this case we denote by $\tilde{H}_k^T$ restriction, to subcollection $H_k^{\mathcal{B}}$ of collection of morphisms $H_\infty^{\mathcal{B}}$ of category of morphisms $V_\infty^{\mathcal{B}}$ of $Q_{\mathcal{B}}$, of k-th horizontal function H_k^T associated to T. Thus defined $\tilde{H}_k^T$ is a function from $H_k^{\mathcal{B}}$ to collection H_k^C. Moreover, we denote by $\tilde{V}_k^T$ restriction, to subcategory $V_k^{\mathcal{B}}$ of category of morphisms $V_\infty^{\mathcal{B}}$ of $Q_{\mathcal{B}}$, of k-th vertical functor V_k^T associated to T. Thus defined $\tilde{V}_k^T$ is a functor from category $V_k^{\mathcal{B}}$ to k-th vertical category V_k^C associated to C.

Lemma 4.13. *Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Let $T, L : Q_{\mathcal{B}} \to C$ be double functors from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to C. If functions $\tilde{H}_1^T$ and $\tilde{H}_1^L$ associated to T and L respectively are equal, then for every positive integer k functions $\tilde{H}_k^T$ and $\tilde{H}_k^L$ are equal and functors $\tilde{V}_k^T$ and $\tilde{V}_k^L$ are equal.*

Proof. Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily generated double category. Let $T, L : Q_{\mathcal{B}} \to C$ be double functors from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to C. Let k be a positive integer. Suppose functions $\tilde{H}_1^T$ and $\tilde{H}_1^L$ associated to T and L respectively are equal. We wish to prove, in that case, that functions $\tilde{H}_k^T$ and $\tilde{H}_k^L$ are equal and that functors $\tilde{V}_k^T$ and $\tilde{V}_k^L$ are equal.

We proceed by induction on k. We first prove that functors $\tilde{V}_1^T$ and $\tilde{V}_1^L$ are equal. Observe first that restriction of morphism function of functor $\tilde{V}_1^T$ to collection $H_1^{\mathcal{B}}$ is equal to function $\tilde{H}_1^T$ and that restriction of morphism function of functor $\tilde{V}_1^L$ to collection $H_1^{\mathcal{B}}$ is equal to function $\tilde{H}_1^L$. From this and from the assumption of the lemma it follows that morphisms function of functors $\tilde{V}_1^T$ and $\tilde{V}_1^L$ to collection $H_1^{\mathcal{B}}$ are equal. We conclude, from this, from the fact that collection $H_1^{\mathcal{B}}$ generates category $V_1^{\mathcal{B}}$, and from functoriality of $\tilde{V}_1^T$ and $\tilde{V}_1^L$, that $\tilde{V}_1^T$ and $\tilde{V}_1^L$ are equal.

Let now k be a positive integer strictly greater than one. Suppose that for every positive integer m strictly less than k functions $\tilde{H}_m^T$ and $\tilde{H}_m^L$ are equal and functors $\tilde{V}_m^T$ and $\tilde{V}_m^L$ are equal. We now prove that functions $\tilde{H}_k^T$ and $\tilde{H}_k^L$ are equal. Observe first that restriction of function $\tilde{H}_k^T$ to collection $\mathrm{Hom}_{V_{k-1}^{\mathcal{B}}}$ of morphisms of category $V_{k-1}^{\mathcal{B}}$ is equal to morphism function of functor $\tilde{V}_{k-1}^T$ and that restriction of function $\tilde{H}_k^L$ to collection $\mathrm{Hom}_{V_{k-1}^{\mathcal{B}}}$ of morphisms of category $V_{k-1}^{\mathcal{B}}$ is equal to morphism function of functor $\tilde{V}_{k-1}^L$. From this, from induction hypothesis, and from the fact that both T and L are double functors it follows that functions $\tilde{H}_k^T$ and $\tilde{H}_k^L$ are equal. We now prove that functors $\tilde{V}_k^T$ and $\tilde{V}_k^L$ are equal. Observe again that restriction of morphism function of functor $\tilde{V}_k^T$ to collection $H_k^{\mathcal{B}}$ is equal to function $\tilde{H}_k^T$ and that restriction of morphism function of functor $\tilde{V}_k^L$ to collection $H_k^{\mathcal{B}}$ is equal to function $\tilde{H}_k^L$. From this, from the previous argument, from the fact that collection $H_k^{\mathcal{B}}$ generates category $V_k^{\mathcal{B}}$, and from functoriality of $\tilde{V}_k^T$ and $\tilde{V}_k^L$ it follows that functors $\tilde{V}_k^T$ and $\tilde{V}_k^L$ are equal. This concludes the proof. ■

Given a bifunctor $T : Q_{\mathcal{B}} \to C$ from free globularily generated double category associated to a decorated bicategory $\mathcal{B}$ to a globularily generated double category C, it is a straightforward observation that morphism functor T_1 of T is equal to limit $\varinjlim \tilde{V}_k^T$ in 2-category **Cat** of sequence $\tilde{V}_k^T$. This, together with lemma 4.13 implies the following proposition. We interpret this by saying that a double functor with domain a free globularily generated double category is completely determined by its value on globular morphisms.

Proposition 4.14. *Let $\mathcal{B}$ be a decorated bicategory. Let C be a globularily gnerated double category. Let $T, L : Q_{\mathcal{B}} \to C$ be double functors from globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to C. If functions $\tilde{H}_1^T$ and $\tilde{H}_1^L$ are equal then morphism functors T_1 and L_1 of double functors T and L are equal.*

Uniqueness part of theorem 1 follows directly from the above proposition. We interpreted completeness part of theorem 1 by saying that every globularily generated verticalization of a decorated bicategory $\mathcal{B}$ could be interpreted as a quotient of free globularily generated dobule category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ via canonical projection. We interpret uniqueness part of theorem 1 by saying that in this case the choice of canonical projections as projection is canonical.

5 Free double functor

We introduce the free double functor construction. We associate, to every decorated bifunctor between decorated bicategories, a double functor between corresponding free globularily generated double categories. We call double functors thus constructed free double functors. Methods used in this section will be a combination of generalizations of methods used in constructions performed in sections 3 and 4.

Construction

Proposition 5.1. *Let $\mathcal{B}, \mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. In that case there exists a pair, formed by a sequence of functions $E_k^G : E_k^{\mathcal{B}} \to E_k^{\mathcal{B}'}$, and a sequence of functors $F_k^G : F_k^{\mathcal{B}} \to F_k^{\mathcal{B}'}$, with k running over all positive integers, such that the following conditions are satisfied:*

1. *Let Φ be a globular morphism in $\mathcal{B}$. The image $E_1^G\Phi$ of Φ under function E_1^G is equal to the image $G\Phi$ of Φ under bifunctor G. Moreover, if α is a vertical morphism of $\mathcal{B}$, then image $E_1^G i_\alpha$, of horizontal identity i_α of morphism α under function E_1^G, is equal to horizontal identity $i_{G\alpha}$ of horizontal morphism $G\alpha$ of $\mathcal{B}'$.*

2. *For every pair of positive integers m, k such that m is less than or equal to k, restriction of function E_k^G to collection of morphisms of category*

$F_m^{\mathcal{B}}$ *is equal to morphism function of functor* F_m^G*, and restriction of morphism function of functor* F_k^G *to collection* $E_m^{\mathcal{B}}$ *equals function* E_m^G*.*

3. *The following two squares commute for every positive integer* k





4. *The following two squares commute for every positive integer* k





5. *The following two squares commute for every positive integer* k





6. *The following square commutes for every positive integer* k



Moreover, conditions 1-5 above determine pair of sequences E_k^G and F_k^G.

Proof. Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. We wish to construct a sequence of functions E_k^G from collection $E_k^{\mathcal{B}}$ to collection $E_k^{\mathcal{B}'}$ and a sequence of functors F_k^G from category $F_k^{\mathcal{B}}$ to category $F_k^{\mathcal{B}'}$ with k running over collection of positive integers, such that pair of sequences E_k^G and F_k^G satisfy conditions 1-6 above.

We proceed inductively on k. We begin with the definition of function E_1^G. Denote first by E_0^G function from $\mathbb{G}^{\mathcal{B}}$ to $\mathbb{G}^{\mathcal{B}'}$ defined as follows: Let Φ be an element of $\mathbb{G}^{\mathcal{B}}$. Assume first that Φ is a globular morphism in $\mathcal{B}$. In that case we make $E_0^G\Phi$ to be equal to the image $G\Phi$ of morphism Φ under functor G. Suppose now Φ is equal to horizontal identity i_α of vertical morphism α of $\mathcal{B}$. In that case we make $E_1^G i_\alpha$ to be equal to horizontal identity $i_{G^*\alpha}$ of the image $G^*\alpha$ of vertical morphism α under decoration G^* of G. A direct computation proves that thus defined, function E_0^G satisfies the following conditions:

1. The following two squares commute



2. The following two squares commute



We now define function $E_1^G : E_1^{\mathcal{B}} \to E_1^{\mathcal{B}'}$ as function $\mu_{E_0^G, G^*}$ associated to pair formed by E_0^G and morphism function of G^*. Function E_1^G is well defined by condition 1 above. By the way it was defined, it is immediate that E_1^G satisfies conditions 1 and 4 in the proposition. Moreover, function E_1^G satisfies condition 6 of the proposition. Finally, condition 2 above implies that function E_1^G satisfies condition 3 of the proposition. We define functor F_1^G as follows: Object function of F_1^G will be restriction, to collection of horizontal morphisms of $\mathcal{B}$, of bifunctor adjacent to G. From the fact that function E_1^G satisfies condition 3 in the proposition it follows that function E_1^G, defined in collection $E_1^{\mathcal{B}}$ of generators of category $F_1^{\mathcal{B}}$, admits a unique extension to a functor from category $F_1^{\mathcal{B}}$ to category $F_1^{\mathcal{B}'}$. We make functor F_1^G to be this extension. It follows, directly from its definition, that functor F_1^G satisfies part 2 of the proposition with respect to function E_1^G. Moreover, the fact that function E_1^G satisfies condition 4 of the proposition implies that functor F_1^G satisfies condition 5 of the proposition.

Let k be a positive integer strictly greater than 1. Assume now that for every positive integer m strictly less than k, a function E_m^G from collection $E_m^{\mathcal{B}}$ to collection $E_m^{\mathcal{B}'}$ and a functor F_m^G from category $F_m^{\mathcal{B}}$ to category $F_m^{\mathcal{B}'}$ have been defined in such a way that pair formed by sequence $E_m^{\mathcal{B}}$ and sequence F_m^G with m running through collection of positive integers strictly less than k, satisfies conditions 1-5 of the first part of the proposition. We now construct a function E_k^G from collection $E_k^{\mathcal{B}}$ to collection $E_k^{\mathcal{B}'}$ and a functor F_k^G from category $F_k^{\mathcal{B}}$ to category $F_k^{\mathcal{B}'}$ such that pair E_k^G and F_k^G satisfies conditions 1-6 above, with respect to pair of sequences E_m^G and F_m^G assumed to exist.

We first define function E_k^G. From the assumption that functor F_{k-1}^G satisfies condition 5 in the proposition with respect to functors s_k and t_k it follows that function $\mu_{F_{k-1}^G, G^*}$ is a well defined function from collection $E_k^{\mathcal{B}}$ to collection $E_k^{\mathcal{B}'}$. We make E_k^G to be this function. From the way it is defined it is clear that function E_k^G satisfies conditions 2,4, and 6 of the proposition. Part 3 of the proposition follows from the functoriality of F_{k-1}^G in category

$F^{\mathcal{B}}_{k-1}$ and from the fact that function E^G_k satisfies part 2 of the proposition. Part 1 of the proposition follows from induction hypothesis. We now define functor F^G_k as follows: We make object function of F^G_k to be restriction of bifunctor F to collection of horizontal morphisms of $\mathcal{B}$. From the fact that function E^G_k satisfies condition 3 of the proposition it follows that E^G_k admits a unique extension to a functor from category $F^{\mathcal{B}}_k$ to category $F^{\mathcal{B}'}_k$. We make F^G_k to be this functor. Thus defined, functor F^G_k satisfies condition 5 of the proposition. This follows from the fact that function E^G_k satisfies condition 4 of the proposition.

We have thus defined, recursively, a pair of sequences E^G_k and F^G_k satisfying conditions 1-6 above. We now prove that these conditions determine pair of sequences E^G_k and F^G_k. Let T_k be a sequence of functions from collection $E^{\mathcal{B}}_k$ to collection $E^{\mathcal{B}'}_k$ and let U_k be a sequence of functors from category $F^{\mathcal{B}}_k$ to category $F^{\mathcal{B}'}_k$ with k running through collection of positive integers. Suppose pair of sequences T_k and U_k satisfy conditions 1-6 above. We prove that for each positive integer k function T_k is equal to function E^G_k and that functor U_k is equal to functor F^G_k. We proceed by induction on k. Observe first that from condition 1 of the proposition both function T_1 and function E^G_1 are equal to function E^G_0 when restricted to collection $\mathbb{G}^{\mathcal{B}}$. From this and from the fact that both T_1 and E^G_1 satisfy conditions 4 and 6 it follows that functions T_1 and E^G_1 are equal. Now, both functor U_1 and F^G_1 are equal on collection of generators $E^{\mathcal{B}}_1$ of $F^{\mathcal{B}}_1$. From this and from the fact that both U_1 and F^G_1 are functors on category $F^{\mathcal{B}}_1$, freely generated by collection $E^{\mathcal{B}}_1$ it follows that functors U_1 and F^G_1 are equal. Let k be a positive integer strictly greater than 1. Suppose now that for every positive integer m strictly less than k functions T_m and E^G_m are equal and functors U_m and F^G_m are equal. From the fact that both function T_k and E^G_k satisfy condition 2 it follows that, when restricted to collection of morphisms of category $F^{\mathcal{B}}_{k-1}$, both functions are equal to morphism function of functor F^G_{k-1}. Now, from the fact that both T_k and E^G_k satisfy conditions 4 and 6 it follows that they are equal. Finally, functors U_k and F^G_k are equal on collection of generators $E^{\mathcal{B}}_k$ of $F^{\mathcal{B}}_k$. From this and from the fact that both U_1 and F^G_1 are functors on category $F^{\mathcal{B}}_k$, freely generated by collection $E^{\mathcal{B}}_k$ it follows that functors U_k and F^G_k are equal. This concludes the proof.

■

Observation 5.2. Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. Let m, k be positive integers such that m is less than or equal to k. In that case condition 2 of proposition above implies that restriction of function E^G_k to collection $E^{\mathcal{B}}_k$ is equal to function E^G_m and

that restriction of functor F_k^G to category $F_m^{\mathcal{B}}$, is equal to functor F_m^G.

Given a decorated bifunctor $G : \mathcal{B} \to \mathcal{B}'$ from a decorated bicategory $\mathcal{B}$ to a decorated bicategory $\mathcal{B}'$ we will write E_∞^G for the limit $\varinjlim E_k^G$ in category **Set** of sequence of functions E_k^G. Thus defined E_∞^G is a function from collection $E_\infty^{\mathcal{B}}$ to collection $E_\infty^{\mathcal{B}'}$. Further, we will write F_∞^G for the limit $\varinjlim F_k^G$ in 2-category **Cat** of sequence of functors F_k^G. Thus defined, F_∞^G is a functor from category $F_\infty^{\mathcal{B}}$ to category $F_\infty^{\mathcal{B}'}$. The following lemma follows directly from proposition 1 and observation 2.

Lemma 5.3. *Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. Function E_∞^G and functor F_∞^G associated to G satisfy the following conditions:*

1. *Function E_∞^G is equal to morphism function of functor F_∞^G.*
2. *Let k be a positive integer. Restriction of function E_∞^G to collection $E_k^{\mathcal{B}}$ is equal to function E_k^G. Moreover, restriction of functor F_∞^G to category $F_k^{\mathcal{B}}$ is equal to functor F_k^G.*
3. *The following squares commute*

$$\begin{array}{ccc} F_\infty^{\mathcal{B}} & \xrightarrow{F_\infty^G} & F_\infty^{\mathcal{B}'} \\ {\scriptstyle s_\infty^{\mathcal{B}}}\downarrow & & \downarrow{\scriptstyle s_\infty^{\mathcal{B}'}} \\ \mathcal{B}^* & \xrightarrow{G^*} & \mathcal{B}'^* \end{array} \qquad \begin{array}{ccc} F_\infty^{\mathcal{B}} & \xrightarrow{F_\infty^G} & F_\infty^{\mathcal{B}'} \\ {\scriptstyle t_\infty^{\mathcal{B}}}\downarrow & & \downarrow{\scriptstyle t_\infty^{\mathcal{B}'}} \\ \mathcal{B}^* & \xrightarrow{G^*} & \mathcal{B}'^* \end{array}$$

4. *The following square commutes*

$$\begin{array}{ccc} E_\infty^{\mathcal{B}} \times_{Hom_{\mathcal{B}^*}} E_\infty^{\mathcal{B}} & \xrightarrow{E_\infty^G \times_G E_\infty^G} & E_\infty^{\mathcal{B}'} \times_{Hom_{\mathcal{B}'^*}} E_\infty^{\mathcal{B}'} \\ {\scriptstyle *_\infty^{\mathcal{B}}}\downarrow & & \downarrow{\scriptstyle *_\infty^{\mathcal{B}'}} \\ E_\infty^{\mathcal{B}} & \xrightarrow{E_\infty^G} & E_\infty^{\mathcal{B}'} \end{array}$$

Lemma 5.4. *Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. Functor F^G_∞ associated to decorated bifunctor G is compatible with equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$ defined on categories $F^{\mathcal{B}}_\infty$ and $F^{\mathcal{B}'}_\infty$ respectively.*

Proof. Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. We wish to prove that functor F^G_∞ associated to G is compatible with equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$ defined on categories $F^{\mathcal{B}}_\infty$ and $F^{\mathcal{B}'}_\infty$ respectively. Compatibility of functor F^G_∞ with relation 1 in the definition of equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$ follows directly from conditions 5 and 6 in proposition 5.1.

We prove now that functor F^G_∞ is compatible with respect to relation 2 in the definition of $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$. Let Φ and Ψ first be globular morphisms in $\mathcal{B}$. Assume that pair Φ, Ψ is composable in $\mathcal{B}$. In that case the image $F^G_\infty\Psi \bullet^{\mathcal{B}}_\infty \Phi$ of vertical composition $\Psi \bullet^{\mathcal{B}}_\infty \Phi$ under functor F^G_∞, is equal to vertical composition $F^G_\infty\Psi \bullet^{\mathcal{B}'}_\infty F^G_\infty\Phi$ in $\mathcal{B}'$ of $F^G_\infty\Phi$ and $F^G_\infty\Psi$. This is, due to condition 2 in proposition 5.1 equal to vertical composition $F^G_1\Psi \bullet^{\mathcal{B}'}_1 F^G_1\Phi$ in $\mathcal{B}'$ of $F^G_1\Phi$ and $F^G_1\Psi$, which in turn is equal to composition $G\Psi \bullet^{\mathcal{B}'}_1 G\Phi$ due to condition 1 of proposition 5.1. Now, the image $F^G_\infty\Psi\Phi$, under functor F^G_∞ of vertical composition $\Psi\Phi$ in $\mathcal{B}$, of globular morphisms Φ and Ψ, is equal, by condition 1 in proposition 5.1 to the image $G\Psi\Phi$ of composition $\Psi\Phi$ under functor G, which in turn is equal to composition $G\Psi G\Phi$ of $G\Phi$ and $G\Psi$. This proves that functor F^G_∞ is compatible with respect to relation 1 in the definition of equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$ when restricted to globular morphisms of $\mathcal{B}$. We now prove functor F^G_∞ is compatible with respect to relation 2 in the definition of $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$ when restricted to horizontal identities of vertical morphisms of $\mathcal{B}$. Let α and β be vertical morphisms of $\mathcal{B}$. Assume that pair α, β is composable in decoration $\mathcal{B}^*$ of $\mathcal{B}$. In that case image $F^G_\infty i_\beta \bullet^{\mathcal{B}}_\infty i_\alpha$ of composition $i_\beta \bullet^{\mathcal{B}}_\infty i_\alpha$ under functor F^G_∞ is equal to composition $F^G_\infty i_\beta \bullet^{\mathcal{B}'}_\infty F^G_\infty i_\alpha$ of $F^G_\infty i_\alpha$ and $F^G_\infty i_\beta$. This in turn is equal, again due to condition 1 of proposition 5.1, to composition $F^G_1 i_\beta \bullet^{\mathcal{B}'}_1 F^G_1 i_\alpha$, which is equal to composition $i_{G^*\beta} \bullet^{\mathcal{B}'}_1 i_{G^*\alpha}$ of $Gi_{G^*\beta}$ and $Gi_{G^*\alpha}$. Now, image $F^G_\infty i_{\beta\alpha}$ of horizontal identity $i_{\beta\alpha}$ of composition $\beta\alpha$ under functor F^G_∞ is, due to conditions 1 and 2 of proposition 5.1 and functoriality of decoration G^* of G, equal to horizontal identity $i_{G^*\beta G^*\alpha}$ of composition $G^*\beta G^*\alpha$ of $G^*\beta$ and $G^*\alpha$. This proves that functor F^G_∞ is compatible with respect to relation 2 in the definition of equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$ when restricted to horizontal identities. We conclude that functor F^G_∞ is compatible with respect to relation 2 in the definition of equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$.

We now prove that functor F^G_∞ is compatible with relation 3 in the definition of equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$. Let Φ and Ψ be globular morphisms in $\mathcal{B}$. Suppose pair Φ, Ψ is compatible with respect to morphism functions of pair of functors $s^{\mathcal{B}}_\infty, t^{\mathcal{B}}_\infty$. In that case the image $F^G_\infty\Psi *^{\mathcal{B}}_\infty \Phi$ of concatenation $\Psi *^{\mathcal{B}}_\infty \Phi$ under functor F^G_∞ is equal, by condition 4 of lemma 5.3 to image $F^G_1\Psi *^{\mathcal{B}}_1 \Phi$ of concatenation $\Psi *^{\mathcal{B}}_1 \Phi$ under functor F^G_1. This in turn is equal, by condition 2 of lemma 5.3, to concatenation $F^G_1\Psi *^{\mathcal{B}'}_1 F^G_1\Phi$ of $F^G_1\Phi$ and $F^G_1\Psi$. Now, the image $F^G_\infty\Psi * \Phi$ of horizontal composition in $\mathcal{B}$, $\Psi * \Phi$, of Φ and Ψ under functor $F^{\mathcal{B}}_\infty$, is equal, by condition 2 of proposition 5.3 and condition 1 to the image $E^G_1\Psi * \Phi$ of horizontal composition $\Psi * \Phi$ under function E^G_1. This is equal to the image $G\Psi * \Phi$ of $\Psi * \Phi$ under function G, which in turn is equal to horizontal composition, in $\mathcal{B}'$, $G\Psi * G\Phi$ of $G\Psi$ and $G\Phi$. This proves that functor F^G_∞ is compatible with relation 3 in the definition of equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$.

Finally, compatibility of functor F^G_∞ with relations 4 and 5 in the definition of relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$ follows from condition 2 of lemma 5.3 and from the fact that bifunctor adjacent to decorated bifunctor G carries left and right identity transformations to left and right identity transformations and associators to associators. This concludes the proof.

■

Given a decorated bifunctor $G : \mathcal{B} \to \mathcal{B}'$ between decorated bicategories $\mathcal{B}$ and $\mathcal{B}'$ we will write V^G_∞ for functor from category $V^{\mathcal{B}}_\infty$ to category $V^{\mathcal{B}'}_\infty$ induced by functor F^G_∞ and equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$. We will write H^G_∞ for morphism function of functor V^G_∞. Thus defined H^G_∞ is function from $H^{\mathcal{B}}_\infty$ to $H^{\mathcal{B}'}_\infty$ induced by function E^G_∞ and equivalence relations $R^{\mathcal{B}}_\infty$ and $R^{\mathcal{B}'}_\infty$. The following proposition follows directly from lemma 5.3.

Proposition 5.5. *Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. Functor V^G_∞ associated to G satisfies the following conditions:*

1. *The following two squares commute*

$$\begin{array}{ccc} V^{\mathcal{B}}_\infty & \xrightarrow{V^G_\infty} & V^{\mathcal{B}'}_\infty \\ {\scriptstyle s^{\mathcal{B}}_\infty}\downarrow & & \downarrow{\scriptstyle s^{\mathcal{B}'}_\infty} \\ \mathcal{B}^* & \xrightarrow{G^*} & \mathcal{B}'^* \end{array} \qquad \begin{array}{ccc} V^{\mathcal{B}}_\infty & \xrightarrow{V^G_\infty} & V^{\mathcal{B}'}_\infty \\ {\scriptstyle t^{\mathcal{B}}_\infty}\downarrow & & \downarrow{\scriptstyle t^{\mathcal{B}'}_\infty} \\ \mathcal{B}^* & \xrightarrow{G^*} & \mathcal{B}'^* \end{array}$$

2. The following square commutes



Notation 5.6. Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. We write Q_G for pair (G^*, V_∞^G) formed by decoration G^* of G and functor V_∞^G associated to G.

The following is the main theorem of this section.

Theorem 5.7. *Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. In that case pair Q_G is a double functor from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to free globularily generated double category $Q_{\mathcal{B}'}$ associated to $\mathcal{B}'$. Restriction, to decorated double category $\mathcal{B}$, of decorated horizontalization H^*Q_G of double functor Q_G associated to G, is equal to decorated bifunctor G.*

Proof. Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor. We wish to prove that in that case pair Q_G is a double functor from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to free globularily generated double category $Q_{\mathcal{B}'}$ associated to $\mathcal{B}'$. Moreover, we wish to prove that restriction to $\mathcal{B}$ of decorated horizontalization H^*Q_G of double functor Q_G associated to G is equal to G.

A direct computation proves that pair Q_G associated to decorated bifunctor G intertwines horizontal identity functors $i_\infty^{\mathcal{B}}$ and $i_\infty^{\mathcal{B}'}$ of $\mathcal{B}$ and $\mathcal{B}'$ respectively. This, together with a direct application of proposition 5.5 implies that pair Q_G associated to G is a double functor. Object function of morphism functor V_∞^G is equal restriction to collection of horizontal morphisms $\mathcal{B}_1$ of $\mathcal{B}$ and restriction of functor V_∞^G to collection of globular 2-morphisms of $\mathcal{B}$ is equal to 2-cell function of decorated bifunctor G. This together with the fact that object functor of double functor Q_G is decoration G^* of G implies that restriction to $\mathcal{B}$ of decorated horizontalization H^*Q_G of double functor Q_G is equal to decorated bifunctor G. This concludes the proof. ■

Definition 5.8. Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$. In that case we call double functor Q_G the free double functor associated to G.

Uniqueness

We now prove that condition appearing in theorem 1 characterizes the condition of a double functor being free. We begin with the following lemma.

Lemma 5.9. *Let $\mathcal{B}, \mathcal{B}'$ be decorated double categories. Let $T : Q_{\mathcal{B}} \to Q_{\mathcal{B}'}$ be a double functor from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to free globularily generated double category $Q_{\mathcal{B}'}$ associated to $\mathcal{B}'$. If restriction to $\mathcal{B}$ of decorated horizontalization H^*T of T defines a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$ then for every positive integer k morphism function of morphism functor of T restricts to a function from $H_k^{\mathcal{B}}$ to $H_k^{\mathcal{B}'}$ and morphism functor of double functor T restricts to a functor from category $V_k^{\mathcal{B}}$ to category $V_k^{\mathcal{B}'}$.*

Proof. Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $T : Q_{\mathcal{B}} \to Q_{\mathcal{B}'}$ be a double functor from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to free globularily generated double category $Q_{\mathcal{B}'}$ associated to $\mathcal{B}'$. Suppose that restriction, to decorated bicategory $\mathcal{B}$, of decorated horizontalization H^*T of double functor T defines a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$. We wish to prove that in this case for every positive integer k morphism function of morphism functor of T restricts to a function from collection $H_k^{\mathcal{B}}$ to collection $H_k^{\mathcal{B}'}$ and that morphism functor of T restricts to a functor from category $V_k^{\mathcal{B}}$ to category $V_k^{\mathcal{B}'}$.

Let first Φ be an element of collection $\mathbb{G}^{\mathcal{B}}$. If Φ is a globular morphism in $\mathcal{B}$ then, by the conditions of the lemma the image $T\Phi$ of Φ under T is a globular morphism in $\mathcal{B}'$ and thus an element of collection $\mathbb{G}^{\mathcal{B}'}$. Let now α be a vertical morphism in $\mathcal{B}$. The fact that T is a double functor implies that the image Ti_α of horizontal identity i_α under T is the horizontal identity of a vertical morphism in $\mathcal{B}'$ and thus is an element of collection $\mathbb{G}^{\mathcal{B}'}$. This, together with the fact that double functor T intertwines source and target functors $s_\infty^{\mathcal{B}}, t_\infty^{\mathcal{B}}$ of $Q_{\mathcal{B}}$ and source and target functors $s_\infty^{\mathcal{B}'}, t_\infty^{\mathcal{B}'}$ of $Q_{\mathcal{B}'}$ implies that the image of collection $H_1^{\mathcal{B}}$ under morphism function of morphism functor T is contained in collection $H_1^{\mathcal{B}'}$. Now, the fact that the image of morphism function of morphism functor of T restricts to a function from collection of generators $H_1^{\mathcal{B}}$ of category $V_1^{\mathcal{B}}$ to collection of generators $H_1^{\mathcal{B}'}$ of category $V_1^{\mathcal{B}'}$, together with functoriality of morphism

functor of double functor T implies that morphism functor of T restricts to a functor from category $V_1^{\mathcal{B}}$ to category $V_1^{\mathcal{B}'}$. An analogous argument proves, inductively on k, that for every positive integer k morphism function of morphism functor of T restricts to a function from collection $H_k^{\mathcal{B}}$ to collection $H_k^{\mathcal{B}'}$ and that morphism functor of double functor T restricts to a functor from category $V_k^{\mathcal{B}}$ to category $V_k^{\mathcal{B}'}$. This concludes the proof. ■

Notation 5.10. Let $\mathcal{B}, \mathcal{B}'$ be decorated double categories. Let $T : Q_{\mathcal{B}} \to Q_{\mathcal{B}'}$ be a double functor from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to free globularily generated double category $Q_{\mathcal{B}'}$ associated to $\mathcal{B}'$. Suppose restriction, to $\mathcal{B}$, of decorated horizontalization H^*T of T defines a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$. Given a positive integer k we write $T_{k,H}$ for restriction to $H_k^{\mathcal{B}}$, of morphism function of morphism functor of T, and we write $T_{k,V}$ for restriction, to category $V_k^{\mathcal{B}}$, of morphism functor of T. We call function $T_{k,H}$ and functor $T_{k,V}$ the k-th horizontal function and the k-th vertical functor associated to T respectively.

Observation 5.11. Let $\mathcal{B}, \mathcal{B}'$ be decorated bicategories. Let $T : Q_{\mathcal{B}} \to Q_{\mathcal{B}'}$ be a double functor from $Q_{\mathcal{B}}$ to $Q_{\mathcal{B}'}$. Suppose restriction to $\mathcal{B}$ of decorated horizontalization H^*T of T defines a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$. In that case morphism functor T_1 of T is equal to limit $\varinjlim T_{k,V}$ in 2-category **Cat** of sequence $T_{k,V}$. In particular, morphism function of morphism functor T_1 of T is equal to limit $\varinjlim T_{k,H}$ in **Set** of sequence of functions $T_{k,H}$.

Given a double functor $T : Q_{\mathcal{B}} \to Q_{\mathcal{B}'}$ from free globularily generated double category $Q_{\mathcal{B}}$ associated to a decorated bicategory $\mathcal{B}$ to free globularily generated double category $Q_{\mathcal{B}'}$ associated to a decorated bicategory $\mathcal{B}'$ such that restriction to $\mathcal{B}$ of decorated horizontalization H^*T of T defines a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$ we call sequence of functions $T_{k,H}$ the free horizontal filtration associated to T and we call sequence of functors $T_{k,V}$ the free vertical filtration associated to T. Horizontal and free horizontal, and vertical and free vertical filtrations of double functor T need not be equal. Observe nevertheless that for every positive integer k restriction to k-th term $H^{Q_{\mathcal{B}}}$ of horizontal filtration of $Q_{\mathcal{B}}$ of k-th term $T_{k,H}$ of free horizontal filtration associated to T is equal to k-th horizontal function H^T associated to T and that restriction to k-th term of vertical filtration $V_k^{Q_{\mathcal{B}}}$ of $Q_{\mathcal{B}}$ of k-th term of of free vertical filtration $T_{k,V}$ associated to T is equal to k-th vertical functor V_k^T associated to T. The following lemma follows directly from lemma 5.9.

Lemma 5.12. *Let $\mathcal{B}, \mathcal{B}'$ be decorated bicategories. Let $T : Q_{\mathcal{B}} \to Q_{\mathcal{B}'}$ be a decorated bifunctor from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to free globularily generated double category $Q_{\mathcal{B}'}$ associated to $\mathcal{B}'$. Let k be a positive integer. Suppose restriction to $\mathcal{B}$ of decorated horizontalization H^*T of T defines a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$. Function $T_{k,H}$ and functor $T_{k,V}$ associated to T satisfy the following conditions:*

1. *Let m be a positive integer less than or equal to k. Restriction to collection of morphisms of category $V_m^{\mathcal{B}}$, of function $T_{k,H}$ is equal to morphism function of functor $T_{k,V}$ and restriction to collection $H_m^{\mathcal{B}}$ of morphism function of functor $T_{k,V}$ is equal to function $T_{k,H}$. Restriction of function $T_{k,H}$ to collection $H_m^{\mathcal{B}}$ is equal to function $T_{m,H}$ and restriction to category $V_m^{\mathcal{B}}$ of functor $T_{k,V}$ is equal to functor $T_{k,V}$.*

2. *The following two squares commute*

$$\begin{array}{ccc} H_k^{\mathcal{B}} & \xrightarrow{T_{k,H}} & H_k^{\mathcal{B}'} \\ {\scriptstyle s_k^{\mathcal{B}}}\downarrow & & \downarrow{\scriptstyle s_k^{\mathcal{B}'}} \\ Hom_{\mathcal{B}^*} & \xrightarrow{T_0} & Hom_{\mathcal{B}'^*} \end{array} \qquad \begin{array}{ccc} H_k^{\mathcal{B}} & \xrightarrow{T_{k,H}} & H_k^{\mathcal{B}'} \\ {\scriptstyle t_k^{\mathcal{B}}}\downarrow & & \downarrow{\scriptstyle t_k^{\mathcal{B}'}} \\ Hom_{\mathcal{B}^*} & \xrightarrow{T_0} & Hom_{\mathcal{B}'^*} \end{array}$$

3. *The following two squares commute*

$$\begin{array}{ccc} H_k^{\mathcal{B}} & \xrightarrow{T_{k,H}} & H_k^{\mathcal{B}'} \\ {\scriptstyle d_k^{\mathcal{B}}}\downarrow & & \downarrow{\scriptstyle d_k^{\mathcal{B}'}} \\ \mathcal{B}_1 & \xrightarrow{T} & \mathcal{B}_1' \end{array} \qquad \begin{array}{ccc} H_k^{\mathcal{B}} & \xrightarrow{T_{k,H}} & H_k^{\mathcal{B}'} \\ {\scriptstyle c_k^{\mathcal{B}}}\downarrow & & \downarrow{\scriptstyle c_k^{\mathcal{B}'}} \\ \mathcal{B}_1 & \xrightarrow{T} & \mathcal{B}_1' \end{array}$$

4. *The following square commutes*

$$\begin{array}{ccc} H_k^{\mathcal{B}} \times_{Hom_{\mathcal{B}^*}} H_k^{\mathcal{B}} & \xrightarrow{T_{k,H} \times_T T_{k,H}} & H_k^{\mathcal{B}'} \times_{Hom_{\mathcal{B}'^*}} H_k^{\mathcal{B}'} \\ \downarrow{\scriptstyle *_k^{\mathcal{B}}} & & \downarrow{\scriptstyle *_k^{\mathcal{B}'}} \\ H_k^{\mathcal{B}} & \xrightarrow{T_{k,H}} & H_k^{\mathcal{B}'} \end{array}$$

5. *The following two squares commute*

$$\begin{array}{ccc} V_k^{\mathcal{B}} & \xrightarrow{T_{k,V}} & V_k^{\mathcal{B}'} \\ \downarrow{\scriptstyle s_{k+1}^{\mathcal{B}}} & & \downarrow{\scriptstyle s_{k+1}^{\mathcal{B}'}} \\ \mathcal{B}^* & \xrightarrow{T_0} & \mathcal{B}'^* \end{array} \qquad \begin{array}{ccc} V_k^{\mathcal{B}} & \xrightarrow{T_{k,V}} & V_k^{\mathcal{B}'} \\ \downarrow{\scriptstyle t_{k+1}^{\mathcal{B}}} & & \downarrow{\scriptstyle t_{k+1}^{\mathcal{B}'}} \\ \mathcal{B}^* & \xrightarrow{T_0} & \mathcal{B}'^* \end{array}$$

The following proposition says that both horizontal and vertical free filtrations of a double functor between free globularily generated double categories are completely determined by their first term.

Proposition 5.13. *Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let T and T' be double functors from $Q_{\mathcal{B}}$ to $Q_{\mathcal{B}'}$. Suppose decorated horizontalizations H^*T and H^*T' of T and T' restrict to decorated bifunctors from $\mathcal{B}$ to $\mathcal{B}'$. If functions $T_{1,H}$ and $T'_{1,H}$ are equal then for every positive integer k functions $T_{k,H}$ and $T'_{k,H}$ are equal and functors $T_{k,V}$ and $T'_{k,V}$ are equal.*

Proof. Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let T and T' be double functors from $Q_{\mathcal{B}}$ to $Q_{\mathcal{B}'}$. Suppose restriction to $\mathcal{B}$ of decorated horizontalizations H^*T and H^*T' of T and T' define decorated bifunctors from $\mathcal{B}$ to $\mathcal{B}'$. Further, assume that functions $T_{1,H}$ and $T'_{1,H}$ are equal. We wish to prove in that for every positive integer k functions $T_{k,H}$ and $T'_{k,H}$ are equal and that functors $T_{k,V}$ and $T'_{k,V}$ are equal.

We proceed by induction on k. We are assuming that functions $T_{1,H}$ and $T'_{1,H}$ are equal. We prove that functors $T_{1,V}$ and $T'_{1,V}$ are equal. Observe that restriction to collection $H_1^{\mathcal{B}}$ of morphism function of functor $T_{1,V}$ is equal to function $T_{1,H}$ and that restriction to collection $H_1^{\mathcal{B}}$ of morphism function of

functor $T'_{1,V}$ is equal to function $T'_{1,H}$. This, the assumption that functions $T_{1,H}$ and $T'_{1,H}$ are equal, the fact that collection $H_1^{\mathcal{B}}$ generates category $V_1^{\mathcal{B}}$, and functoriality of $T_{1,V}$ and $T'_{1,V}$ implies that functors $T_{1,V}$ and $T'_{1,V}$ are equal.

Let k be a positive integer strictly greater than 1. Suppose that for every positive integer m strictly less than k functions $T_{m,H}$ and $T'_{m,H}$ are equal and functors $T_{m,V}$ and $T'_{m,V}$ are equal. We now prove that functions $T_{k,H}$ and $T'_{k,H}$ are equal and that functors $T_{k,V}$ and $T'_{k,V}$ are equal. We first prove that functions $T_{k,H}$ and $T'_{k,H}$ are equal. Morphism functions of functors $T_{k-1,V}$ and $T'_{k-1,V}$ are equal by induction hypothesis. This and the fact that functors $T_{k-1,V}$ and $T'_{k-1,V}$ both satisfy conditions 4 and 5 of lemma 5.12 implies that functions $T_{k,H}$ and $T'_{k,H}$ are equal. We now prove that functors $T_{k,V}$ and $T'_{k,V}$ are equal. Restriction to collection $H_k^{\mathcal{B}}$ of morphism function of functor $T_{k,V}$ is equal to function $T_{k,H}$ and restriction to collection $H_k^{\mathcal{B}}$ of morphism function of functor $T'_{k,V}$ is equal to function $T'_{k,H}$. The fact that functions $T_{k,H}$ and $T'_{k,H}$ are equal, the fact that collection $H_k^{\mathcal{B}}$ generates category $V_k^{\mathcal{B}}$, and functoriality of both $T_{k,V}$ and $T'_{k,V}$ implies that functors $T_{k,V}$ and $T'_{k,V}$ are equal. This concludes the proof. ■

Corollary 5.14. *Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let T and T' be double functors from $Q_{\mathcal{B}}$ to $Q_{\mathcal{B}'}$. Suppose decorated horizontalizations H^*T and H^*T' of T and T' restrict to decorated bifunctors from $\mathcal{B}$ to $\mathcal{B}'$. If functions $T_{1,H}$ and $T'_{1,H}$ are equal then T and T' are equal.*

We conclude this section with the following theorem, which says that conditions in theorem 1 characterize the condition of a double functor being free. The proof of this theorem is a direct consequence of corollary 5.14.

Theorem 5.15. *Let $\mathcal{B}$ and $\mathcal{B}'$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ be a decorated bifunctor from $\mathcal{B}$ to $\mathcal{B}'$. Double functor Q_G is the only double functor from free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to free globularily generated double category $Q_{\mathcal{B}'}$ associated to $\mathcal{B}'$ such that the restriction, to $\mathcal{B}$, of its decorated horizontalization, is equal to G.*

6 Freeness

We prove that globularily generated double category construction presented in section 3 and free double functor construction presented in section 5 combine into a functor. We call this functor the free globularily generated double category functor. We prove that free globularily generated double cateogry

functor is left adjoint to decorated horizontalization functor. We interpret this result by saying that free globularily generated double category construction is free with respect to decorated horizontalization functor.

Functoriality

We begin by proving that free double functor construction presented in section 5 provides free globularily generated double category construction presented in section 3 with the structure of a functor. We begin with the following lemma.

Lemma 6.1. *Let $\mathcal{B}, \mathcal{B}'$, and $\mathcal{B}''$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ and $G' : \mathcal{B}' \to \mathcal{B}''$ be decorated bifunctors. Let k be a positive integer. In that case function $H_k^{G'G}$ associated to composition $G'G$ is equal to composition $H_k^{G'}H_k^G$ of functions H_k^G and $H_k^{G'}$, and functor $V_k^{G'G}$ associated to composition $G'G$ is equal to composition $V_k^{G'}V_k^G$ of functors V_k^G and $V_k^{G'}$.*

Proof. Let $\mathcal{B}, \mathcal{B}'$, and $\mathcal{B}''$ be decorated bicategories. Let $G : \mathcal{B} \to \mathcal{B}'$ and $G' : \mathcal{B}' \to \mathcal{B}''$ be decorated bifunctors. Let k be a positive integer. We wish to prove that in this case function $H_k^{G'G}$ associated to composition $G'G$ of G and G' is equal to composition $H_k^{G'}H_k^G$ of functions H_k^G and $H_k^{G'}$. Moreover, we wish to prove that functor $V_k^{G'G}$ associated to composition $G'G$ of G and G' is equal to composition $V_k^{G'}V_k^G$ of functors V_k^G and $V_k^{G'}$.

We proceed by induction on k. We first prove that function $H_1^{G'G}$ associated to composition $G'G$ of G and G' is equal to composition $H_1^{G'}H_1^G$ of H_1^G and $H_1^{G'}$. Let Φ be an element of collection $\mathbb{G}^{\mathcal{B}}$. Suppose first that Φ is a globular morphism in $\mathcal{B}$. In that case the image $H_1^{G'G}\Phi$ of Φ under function $H_1^{G'G}$ is equal to the image $G'G\Phi$ of Φ under composition $G'G$ of G and G', this on the other hand is equal to composition $G'\Phi G\Phi$ of globular morphisms $G\Phi$ and $G'\Phi$ in $\mathcal{B}'$. Now, the image $H_1^{G'}H_1^G\Phi$ of Φ under composition $H_1^{G'}H_1^G$ is equal to the image $H_1^{G'}H_1^G\Phi$ of $H_1^G\Phi$ under function $H_1^{G'}$. The image $H_1^G\Phi$ of Φ under function H_1^G is equal to the image $G\Phi$ of Φ under G. Now, the image $H_1^{G'}G\Phi$ of globular morphism $G\Phi$ in $\mathcal{B}'$ under function $H_1^{\mathcal{B}'}$ is equal to the image $G'G\Phi$ of morphism $G\Phi$ under G', which in turn is equal to the image $G'G\Phi$ of Φ under composition $G'G$ of G and G'. This proves that function $H_1^{G'G}$ associated to composition $G'G$ is equal to composition $H_1^{G'}H_1^G$ when restricted to globular morphisms of $\mathcal{B}$. Suppose now that Φ is the horizontal identity i_α of a vertical morphism α of $\mathcal{B}$. In this case the image $H_1^{G'G}\Phi$ of Φ under function $H_1^{G'G}$ is equal to the image $G'G\Phi$ of Φ under composition $G'G$ of G and G', which is equal to horizontal

identity $i_{G'G\alpha}$ of vertical morphism $G'G\alpha$ in $\mathcal{B}''$. Now, the image $H_1^{G'}H_1^G\Phi$ of Φ under composition $H_1^{G'}H_1^G$ is equal to the image of $H_1^G\Phi$ under function $H_1^{G'}$. Now, the image $H_1^G\Phi$ of Φ under H_1^G is equal to horizontal identity $i_{G\alpha}$ of vertical morphism $G\alpha$ in $\mathcal{B}'$. The image $H_1^{G'}H_1^G\Phi$ of $H_1^G\Phi$ under function $H_1^{G'}$ is thus the image of horizontal identity $i_{G\alpha}$ under $H_1^{G'}$, which in turn is equal to horizontal identity $i_{G'G\alpha}$ of vertical morphism $G'G\alpha$ in $\mathcal{B}''$. This proves that function $H_1^{G'G}$ is equal to composition $H_1^{G'}H_1^G$ when restricted to collection $\mathbb{G}^{\mathcal{B}}$. This and the fact that functions $H_1^G, H_1^{G'}$, and $H_1^{G'G}$ all satisfy condition 2 of proposition 5.5 proves that equality extends to collection $H_1^{\mathcal{B}}$. Now both functor $V_1^{G'G}$ associated to composition $G'G$ and composition $V_1^{G'}V_1^G$ of functors $V_1^{G'}$ and V_1^G are equal to function $H_1^{G'G}$ when restricted to collection $H_1^{\mathcal{B}}$ of generators of category $V_1^{\mathcal{B}}$. This and the fact that function $H_1^{G'G}$ satisfies condition 2 of proposition 5.5 implies that functor $V_1^{G'G}$ is equal to composition $V_1^{G'}V_1^G$ of $V_1^{G'}$ and V_1^G.

Let now k be a positive integer strictly greater than 1. Suppose that for every positive integer m strictly less than k equations $H_m^{G'G} = H_m^{G'}H_m^G$ and $V_m^{G'G} = V_m^{G'}V_m^G$ are satisfied. We now prove that equations $H_k^{G'G} = H_k^{G'}H_k^G$ and $V_k^{G'G} = V_k^{G'}V_k^G$ are satisfied. We first prove that function $H_k^{G'G}$ associated to composition $G'G$ of G and G' is equal to composition $H_k^{G'}H_k^G$ of H_k^G and $H_k^{G'}$. Both function $H_k^{G'G}$ associated to $G'G$ and composition $H_k^{G'}H_k^G$ of H_k^G and $H_k^{G'}$ are equal to morphism function of functor $V_{k-1}^{G'G}$ when restricted to collection of morphisms of category $V_{k-1}^{G'G}$. This, together with the way function $H_k^{G'G}$ is defined, and the fact that functor $V_{k-1}^{G'G}$ satisfies condition ? of proposition ? implies that function $H_k^{G'G}$ and composition $H_k^{G'}H_k^G$ of $H_k^{G'}$ and H_k^G are equal. Now, both functor $V_k^{G'G}$ associated to composition $G'G$ of G and G' and composition $V_k^{G'}V_k^G$ of functors V_k^G and $V_k^{G'}$ are equal to function $H_k^{G'G}$ when restricted to collection $H_k^{\mathcal{B}}$ of generators of category $V_k^{\mathcal{B}}$. This, together with the fact that function $H_k^{G'G}$ satisfies condition 1 of porposition 5.5 implies that functor $V_k^{G'G}$ associated to composition $G'G$ of G and G' and composition $V_k^{G'}V_k^G$ of V_k^G and $V_k^{G'}$ are equal. This concludes the proof. ■

An argument similar to the one used in lemma 6.1 proves the following lemma.

Lemma 6.2. *Let $\mathcal{B}$ be a decorated bicategory. Let k be a positive integer. In that case function $H_k^{id_{\mathcal{B}}}$ associated to identity decorated biendofunctor $id_{\mathcal{B}}$ of $\mathcal{B}$ is equal to identity function $id_{H_k^{\mathcal{B}}}$ of collection $H_k^{\mathcal{B}}$, and functor $V_k^{id_{\mathcal{B}}}$ associated to identity biendofunctor $id_{\mathcal{B}}$ of $\mathcal{B}$ is equal to endofunctor $id_{V_k^{\mathcal{B}}}$ of category $V_k^{\mathcal{B}}$.*

Notation 6.3. Let k be a positive integer. We write V_k for pair formed by function associating category $V_k^{\mathcal{B}}$ to every decorated bicategory $\mathcal{B}$ and function associating functor V_k^G to every decorated bifunctor G. We write H_k for pair formed by function associating collection $H_k^{\mathcal{B}}$ to every decorated bicategory $\mathcal{B}$ and function associating function H_k^G to every decorated bifunctor G.

Lemma 6.4. *Let k be an integer. Pairs V_k and H_k defined above define functors from category $\boldsymbol{bCat}^*$ of decorated bicategories and decorated bifunctors to categories $\boldsymbol{Cat}$ and $\boldsymbol{Set}$ respectively. Moreover, when restricted to subcategory $\boldsymbol{2Cat}^*$ of $\boldsymbol{bCat}^*$ of decorated 2-categories and decorated bifunctors, functor H_k admits the structure of a functor from $\boldsymbol{2Cat}^*$ to category $\boldsymbol{Cat}$.*

Notation 6.5. We will write V_∞ for pair formed by function associating category $V_\infty^{\mathcal{B}}$ to every decorated bicategory $\mathcal{B}$ and function associating functor V_∞^G to every decorated bifunctor G. We write H_∞ for pair formed by function associating collection of morphisms $H_\infty^{\mathcal{B}}$ of cateogry $V_\infty^{\mathcal{B}}$ to every decorated bicategory $\mathcal{B}$ and morphism function H_∞^G of functor V_∞^G to every decorated bifunctor G.

The following proposition is a direct consequence of lemma 6.4.

Proposition 6.6. *Pairs V_∞ and H_∞ defined above define functors from category $\boldsymbol{bCat}^*$ of decorated bicategories and decorated bifunctors to categories $\boldsymbol{Cat}$ and $\boldsymbol{Set}$ respectively. When restricted to subcategory $\boldsymbol{2Cat}^*$ of $\boldsymbol{bCat}^*$ of decorated 2-categories and decorated bifunctors, functor H_∞ admits the structure of a functor from $\boldsymbol{2Cat}^*$ to category $\boldsymbol{Cat}$. Finally, functors V_∞ and H_∞ are equal to limits $\varinjlim V_k$ and $\varinjlim H_k$ in 2-category $\boldsymbol{Cat}$ of sequences of functors V_k and H_k respectively.*

Proposition 6.7. *Pair formed by function associating free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ to every decorated bicategory $\mathcal{B}$ and function associating free double functor Q_G to every decorated bifunctor G defines a functor from category $\boldsymbol{bCat}^*$ of decorated bicategories and decorated bifunctors, to category $\boldsymbol{gCat}$ of globularily generated double categories and double functors. Further, restriction of functor H_∞ to cateogory $\boldsymbol{2Cat}^*$ of decorated 2-categories and decorated bifunctors is equal to composition τQ of functor Q and restriction to $\boldsymbol{gCat}$ of transversal functor τ.*

Definition 6.8. We call functor Q defined in proposition 6.7 the free globularily generated double category functor.

Adjoint relation

We now prove that free globularily generated functor Q of proposition 6.7 is left adjoint to decorated horizontalization bifunctor H^*. We begin by proving that canonical double projection construction performed in section 4 is natural. We will write π for collection of canonical double projections π^C with C running over collection of globularily generated double categories.

Proposition 6.9. *Collection π of canonical double projections is a natural transformation.*

Proof. We wish to prove that collection π formed by canonical projections π^C with C running through collection of globularily generated double categories, is a natural transformation from composition H^*Q of free globularily generated double category functor Q and decorated horizontalization functor H^* to identity endofunctor $id_{\mathbf{gCat}}$ of category **gCat** of globularily generated double categories and double functors. That is, we wish to prove that for every double functor $T : C \to C'$ from a globularily generated double category C to a globularily generated double category C' the following square commutes

$$\begin{array}{ccc} Q_{H^*C} & \xrightarrow{Q_{H^*T}} & Q_{H^*C'} \\ \downarrow{\scriptstyle \pi^C} & & \downarrow{\scriptstyle \pi^{C'}} \\ C & \xrightarrow{T} & C' \end{array}$$

Let C and C' be globularily generated double categories. Let $T : C \to C'$ be a double functor from C to C'. We first prove that for each positive integer k the following two squares commute

$$\begin{array}{ccc} H_k^{H^*C} & \xrightarrow{H_k^{H^*T}} & H_k^{H^*C'} \\ \downarrow{\scriptstyle H_k^{\pi^C}} & & \downarrow{\scriptstyle H_k^{\pi^{C'}}} \\ \mathrm{Hom}_{C_1} & \xrightarrow{T} & \mathrm{Hom}_{C_1'} \end{array} \qquad \begin{array}{ccc} V_k^{H^*C} & \xrightarrow{V_k^{H^*T}} & V_k^{H^*C'} \\ \downarrow{\scriptstyle V_k^{\pi^C}} & & \downarrow{\scriptstyle V_k^{\pi^{C'}}} \\ C_1 & \xrightarrow{T} & C_1' \end{array}$$

we do this by induction on k. We fist prove that square on the left commutes in the case in which k is equal to 1. Let Φ be an element of collection $\mathbb{G}^{\mathcal{B}}$. Suppose first that Φ is a globular morphism in $\mathcal{B}$. In that case the image $H_1^{H^*T}\Phi$ of Φ under function $H_1^{H^*T}$ is equal to the image $H^*T\Phi$ of Φ under functor H^*T, which is equal to the image $T\Phi$ of Φ under T. The image $H_1^{\pi^{C'}}T\Phi$ of $T\Phi$ under function $H_1^{\pi^{C'}}$ is equal to $T\Phi$. Thus the upper right corner of left side square above, in the case in which k equals 1, is equal to the image $T\Phi$ of Φ under double functor T. Now, the image $H_1^{\pi^C}\Phi$ of Φ under function $H_1^{\pi^C}$ is equal to Φ, which proves that the lower left corner of left square above is also equal to $T\Phi$. Suppose now that Φ is equal to i_α for a vertical morphism α of $\mathcal{B}$. In that case, the image $H_1^{H^*T}\Phi$ of Φ under function $H_1^{H^*T}$ is equal to the image $H^*T\Phi$ of Φ under H^*T, which is, in this case, equal to horizontal identity $i_{H^*T\alpha}$ of vertical morphism $H^*T\alpha$ of C'. This is equal to $T\alpha$ and thus the image of Φ under the upper right corner of the left square above is horizontal identity $i_T\alpha$ of vertical morphism $T\alpha$ in $\mathcal{B}'$. Now, the image $H_1^{\pi^C}$ under function $H_1^{\pi^C}$ of Φ is equal to Φ. Thus the image of Φ under the bottom left corner of diagram on the left above is also equal to $T\Phi$. We conclude that diagram on the left above commutes when restricted to collection $\mathbb{G}^{\mathcal{B}}$. This together with the fact that function $H_1^{H^*T}$ satisfies condition 6 of proposition 5.1 and T is a double functor, implies that square on the left above commutes on collection $H_1^{\mathcal{B}}$. Now, square on the right above commutes when restricted to collection $E_1^{\mathcal{B}}$ of generators of category $V_1^{\mathcal{B}}$. This, together with the fact that all edges involved are functors implies that square on the right above commutes on full category $V_1^{\mathcal{B}}$.

Let now k be a positive integer strictly greater than 1. Suppose that for every positive integer m strictly less than k squares above commute. We now prove that squares above commute for k. Square on the left above commutes when restricted to collection of morphisms of category $V_{k-1}^{H^*C}$ by induction hypothesis. This, together with the fact that the upper edge of the square satisfies condition 4 of proposition 5.12, its left and right edges satisfy condition 5 of 5.12 and its bottom edge satisfies condition ? of definition ? implies that the full square commutes. Now, square on the right above commutes when restricted to collection $E_k^{\mathcal{B}}$ of generators of category $V_k^{\mathcal{B}}$. This, together with the fact that all edges of the square are functors, implies that the full square commutes for category $V_k^{\mathcal{B}}$. The result follows from this by taking limits of squares on the right above.

■

Notation 6.10. Let $\mathcal{B}$ be a decorated double category. In that case $\mathcal{B}$ is a sub decorated double category of decorated horizontalization $H^*Q_{\mathcal{B}}$ of free

globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$. We denote by $j^{\mathcal{B}}$ inclusion decorated bifunctor of $\mathcal{B}$ in $H^*Q_{\mathcal{B}}$. We write j for collection of inclusion decorated bifunctors $j^{\mathcal{B}}$ with $\mathcal{B}$ running through collection of decorated bicategories.

Observation 6.11. As defined above collection of inclusion decorated bifunctors j is clearly a natural transformation.

Theorem 6.12. *Free globularily generated double category functor Q is left adjoint to decorated horizontalization bifunctor H^*. Natural transformations π and j as defined above form a counit-unit pair for adjoint pair Q, H^*.*

Proof. We wish to prove that pair formed by free globularily generated double category functor Q and decorated horizontalization functor H^* forms an adjoint pair. Moreover, we wish to prove that pair formed by canonical projection natural transformation π and natural transformation j forms a counit-unit pair for pair Q, H^*.

It has already been established that π is a natural transformation from composition QH^* of decorated horizontalization functor H^* and free globularily generated double category functor Q to identity endofunctor $id_{\mathbf{gCat}}$ of category **gCat** of globularily generated double categories and double functors, and that j is a natural transformation from identity endofunctor $id_{\mathbf{bCat}^*}$ of category $\mathbf{bCat}^*$ of decorated bicategories and decorated bifunctors, to composition H^*Q of free globularily generated double category functor Q and decorated horizontalization functor H^*. We thus only need to prove that pair π, j satisfies the triangle equations for a counit-unit pair. We begin by proving that the following triangle commutes



Let C be a globularily generated double category. In that case decoration and collection of horizontal morphisms of both decorated horizontalization H^*C of C and decorated horizontalization $H^*Q_{H^*C}$ of free globularily generated double category Q_{H^*C} associated to horizontalization H^*C of C are equal to category of objects C_0 of C and to collection of objects of morphism

category C_1 of C respectively. Moreover, restriction of inclusion decorated bifunctor j_{H^*C} of decorated horizontalization H^*C of C in decorated horizontalization $H^*Q_{H^*C}$ of Q_{H^*C} to both category of objects C_0 of C and collection of horizontal morphisms of C is the identity function. Restriction of inclusion j_{H^*C} to collection of globular morphisms of H^*C is inclusion of collection of globular morphisms of C to collection of globular morphisms of Q_C. Now, again restriction to both decoration and collection of horizontal morphisms of horizontalization $H^*\pi^C$ of canonical projection π^C associated to C is equal to the identity function in both cases. Restriction of $H^*\pi^C$ to collection of globular morphisms of decorated horizontalization $H^*Q_{H^*C}$ of free globularily generated double category Q_{H^*C} associated to decorated horizontalization H^*C of C is equal to restriction, to collection of globular morphisms of C, of canonical projection π^C associated to C, which is equal to the identity function. It follows, from this, that composition $H^*\pi^C j_{H^*C}$ of inclusion j_{H^*C} of decorated horizontalization H^*C associated to C in decorated horizontalization $H^*Q_{H^*C}$ associated to Q_{H^*C} and decorated horizontalization $H^*\pi^C$ of canonical projection π^C associated to C is equal to the identity of decorated horizontalization H^*C. We conclude that triangle above commutes.

We now prove that the following triangle is commutative



Let $\mathcal{B}$ be a decorated bicategory. In this case restriction to both category of objects C_0 of C and to collection of horizontal morphisms of C, of inclusion decorated bifunctor $j_{\mathcal{B}}$ of $\mathcal{B}$ in decorated horizontalization $H^*Q_{\mathcal{B}}$ of free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$ is equal to the identity function. Restriction, to both category of objects C_0 of C and to collection of horizontal morphisms of C, now of canonical projection $\pi_{Q_{\mathcal{B}}}$ associated to free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$, is again equal to the identity function. We now prove that restriction, to decorated bicategory $\mathcal{B}$, of decorated horizontalization $H^*\pi^{Q_{\mathcal{B}}}Q_{j_{\mathcal{B}}}$ of composition $\pi^{Q_{\mathcal{B}}}Q_{j_{\mathcal{B}}}$ of free globularily generated functor $Q_{j_{\mathcal{B}}}$ associated to canonical inclusion $j_{\mathcal{B}}$ associated to $\mathcal{B}$ and canonical projection $\pi^{Q_{\mathcal{B}}}$ associated to free globularily generated double category $Q_{\mathcal{B}}$ associated to $\mathcal{B}$

defines a decorated endobifunctor of $\mathcal{B}$. It has already been established that restriction, to both decoration and collection of horizontal morphisms of $\mathcal{B}$, of both $Q_{j_{\mathcal{B}}}$ and $\pi^{Q_{\mathcal{B}}}$ and thus of composition $\pi^{Q_{\mathcal{B}}}Q_{j_{\mathcal{B}}}$ is equal to the identity function. Now, let Φ be a globular morphism in $\mathcal{B}$ In that case the image $Q_{j_{\mathcal{B}}}\Phi$ of Φ under functor $Q_{j_{\mathcal{B}}}$ is equal to the image $j_{\mathcal{B}}\Phi$ of Φ under canonical inclusion $j_{\mathcal{B}}$ associated to $\mathcal{B}$, which is equal to $\mathcal{B}$. Now, the image $\pi^{Q_{\mathcal{B}}}\Phi$ of Φ under canonical projection $\pi^{Q_{\mathcal{B}}}$ associated to $Q_{\mathcal{B}}$, is again equal to Φ. We conclude that restriction to decorated bicategory $\mathcal{B}$, of decorated horizontalization $H^*\pi^{Q_{\mathcal{B}}}Q_{j_{\mathcal{B}}}$ of composition $\pi^{Q_{\mathcal{B}}}Q_{j_{\mathcal{B}}}$ defines a decorated endobifunctor of $\mathcal{B}$. Moreover, this decorated endobifunctor of $\mathcal{B}$ is the identity endobifunctor of $\mathcal{B}$. It follows, from this, and from proposition 6.9 that composition $\pi^{Q_{\mathcal{B}}}Q_{j_{\mathcal{B}}}$ is equal to the identity endobifunctor of $Q_{\mathcal{B}}$. We conclude that triangle above commutes. This concludes the proof.

■

We interpret theorem 6.12 by saying that free globularily generated double category functor Q is free with respect to decorated horizontalization functor.

7 References

[1] Arthur Bartels, Christopher L. Douglas, Andre Henriques. Dualizability and index of subfactors *Quantum topology* ***5****. 2014. 289-345.*

[2] Christopher L. Douglas, Andre Henriques. Internal bicategories.

[3] Charles Ehresmann. Categories structurees. *Ann. Sci. Ecole Nor. Sup. (3), 80:349-426, 1963.*

[4] Charles Ehresmann. Categories et structures:extraits. *Seminaire Ehresmann. Topologie et geometrie differentielle 6 1-3, 1964.*

[5] Max Kelly, Ross Street. Review of the elements of 2-categories. *Category seminar, Proc. Sem. Sydney, 1972/1973, Lecture notes in math. Vol. 420. 75-103. Springer, Berlin, 1974.*

[6] Saunders Mac Lane. Categories for the working mathematician, *Volume 5, Graduate texts in mathematics. Springer, second edition, 1998.*

[7] Juan Orendain. Globularily generated bicategories I: The globularily generated piece construction.

[8] Michael A. Schulman. Constructing symmetric monoidal bicategories.

[9] N.P. Landsman. Bicategories of operator algebras and possion manifolds.